



Federatie
**Medisch
Specialisten**

Repositiemanoeuvres als behandeling BPPD

Inhoudsopgave

Repositiemanoevres als behandeling BPPD	1
Inhoudsopgave	2
Startpagina - Benigne paroxysmale positieduizeligheid	3
De diagnose van BPPD	5
De diagnose van BPPD van het posterieure kanaal	6
De diagnose van BPPD van het horizontale kanaal	14
De diagnose van BPPD van het anterieure kanaal	19
De differentiaal diagnose van BPPD	23
Het aanvullend onderzoek van BPPD	28
Indicatie voor beeldvormend, audiologisch en vestibulair onderzoek bij BPPD	29
Indicatie voor audiometrisch onderzoek bij BPPD	34
De behandeling van BPPD	38
Repositiemanoevres als behandeling van BPPD	39
Vestibulaire revalidatie als behandeling van BPPD	52
Medicatie als behandeling van BPPD	56
Chirurgische interventie als behandeling van BPPD	60
Expectatief beleid bij BPPD	64
Omgevingsfactoren die belangrijk zijn voor de behandeling van BPPD	68
De herevaluatie van de behandeling van BPPD	71
De noodzaak van herevaluatie van de behandeling van BPPD	72
De inhoud van de herevaluatie van de behandeling van BPPD	77
De voorlichting van patiënten over BPPD	82

Startpagina - Benigne paroxysmale positieduizeligheid

Waar gaat deze richtlijn over?

Deze richtlijn richt zich op wat volgens de huidige maatstaven de beste zorg is voor patiënten met BPPD. In de richtlijn komen de volgende onderwerpen aan de orde:

- Diagnose van BPPD
- Differentiaal diagnose van BPPD
- Aanvullend onderzoek bij BPPD
- Behandeling van BPPD
- Omgevingsfactoren van BPPD
- Herevaluatie van behandeling van BPPD
- Voorlichting patiënten over BPPD

Voor wie is deze richtlijn bedoeld?

Deze richtlijn is bestemd voor alle zorgverleners die betrokken zijn bij de zorg voor patiënten met BPPD.

Voor patiënten

Benigne paroxysmale positieduizeligheid (BPPD) is een kortdurende draaiduizeligheid. De duizeligheid komt op bij bewegingen van het hoofd, bijvoorbeeld bij voorover buigen, omhoog kijken, gaan liggen in bed of omdraaien in bed. De klachten worden veroorzaakt door een soort gruis dat rondzwerft door het evenwichtsorgaan. Het woord paroxysmaal geeft aan dat de duizeligheid in aanvallen optreedt. De klachten houden gewoonlijk niet langer dan één minuut aan. Sommige patiënten zijn gedurende de duizeligheid ook misselijk of moeten braken. BPPD is een onschuldige aandoening die vaak vanzelf weer overgaat. Het woord benigne geeft dat aan.

BPPD is een van de vele soorten van duizeligheid. Ongeveer een op de twintig mensen heeft last van een vorm van duizeligheid. Ongeveer een op de vijf patiënten met duizeligheid heeft BPPD.

Bij aanverwante informatie is een link te vinden naar patiënteninformatie van de KNO-artsen.

Meer informatie over BPPD is ook te vinden op Thuisarts:

<https://www.thuisarts.nl/bppd>

Meer informatie over BPPD is ook te vinden op de website van de neurologen:

<https://www.neurologie.nl/publiek/patientenvoorlichting/bppd>

Hoe is de richtlijn tot stand gekomen?

Het initiatief voor deze richtlijn is afkomstig van Nederlandse Vereniging voor Keel-, Neus- en Oorheelkunde en Heelkunde van het Hoofd-Halsgebied (NVKNO). De richtlijn is opgesteld door een multidisciplinaire commissie met vertegenwoordigers vanuit de KNO-artsen, neurologen en klinische fysici-vestibulogen. In de huidige commissie zijn ook de fysiotherapeuten vertegenwoordigd.

Verantwoording

Laatst beoordeeld : 01-02-2020

Laatst geautoriseerd : 01-02-2020

Voor de volledige verantwoording, evidence tabellen en eventuele aanverwante producten raadpleegt u de Richtlijndatabase.

De diagnose van BPPD

Het onderwerp 'diagnose van BPPD' wordt uitgewerkt in verschillende modules.

Verantwoording

Laatst beoordeeld : 01-02-2020

Laatst geautoriseerd : 01-02-2020

Voor de volledige verantwoording, evidence tabellen en eventuele aanverwante producten raadpleegt u de Richtlijndatabase.

De diagnose van BPPD van het posterieure kanaal

Uitgangsvraag

Wat is de beste manier om BPPD van het posterieure kanaal (p-BPPD) te diagnosticeren?

Aanbeveling

De diagnose BPPD van het posterieur kanaal wordt gesteld wanneer draaiduizeligheid met nystagmus wordt opgewekt door de Dix-Hallpike manoeuvre.

Overwegingen

- Voordeel: duidelijkheid omtrent de diagnose
- Nadelen: het mogelijk provoceren van draaiduizeligheid
- Kosten: minimaal
- Afweging van voordeel tegen nadeel: het voordeel weegt zwaarder.
- Waarde oordelen: Dix-Hallpike manoeuvre is de gouden standaard testmethode voor het stellen van de diagnose BPPD
- Rol van de voorkeur van de patiënt: minimaal.
- Exclusie: patiënten met fysieke beperkingen van de nek, zoals ernstige reumatoïde arthritis en cervicale radiculopathie.




Figure 2.1

Diagrammatic representation of performance of the Dix-Hallpike maneuver for the diagnosis of posterior canal BPPV (adapted from Furman et al., 1999). (A) The examiner stands at the patient's right side and rotates the patient's head 45 degrees to the right to align the right posterior semicircular canal with the sagittal plane of the body. (B) The examiner moves the patient, whose eyes are open, from the seated to the supine right-ear-down position and then extends the patient's neck slightly so that the chin is pointed slightly upward. The latency, duration, and direction of nystagmus, if present, and the latency and duration of vertigo, if present, should be noted. The *arrows* in the inset depict the direction of nystagmus in patients with typical benign paroxysmal positional vertigo. A presumed location in the labyrinth of the free-floating debris thought to cause the disorder is also shown.

Onderbouwing

Conclusies

Niveau 2	De Dix-Hallpike manoeuvre is de gouden standaard om posterieur kanaal BPPD te diagnosticeren.
Niveau 4	<p>Het is niet aangetoond, maar het lijkt waarschijnlijk dat video-oculografie behulpzaam is bij het interpreteren van de nystagmus en biedt het voordeel dat documentatie van de nystagmus mogelijk is. De experts zijn van mening dat voor het beoordelen van de nystagmus een Frenzel bril niet strikt noodzakelijk is.</p> <p><i>Niveau D: bronnen (niet-vergelijkend onderzoek Jackson et al 2007, en case studie Bertholon 2002))</i></p>

Samenvatting literatuur

Posterior semicircular canal BPPV is diagnosed when 1) patients report a history of vertigo provoked by changes in head position relative to the gravity vector and 2) when, on physical examination, characteristic nystagmus is provoked by the Dix-Hallpike maneuver (Table 2.1).

Table 2.1. Diagnostic criteria for posterior canal BPPV

History	Patient reports repeated episodes of vertigo with changes in head position.
Physical examination	<p>Each of the following criteria are fulfilled:</p> <ul style="list-style-type: none"> ● Vertigo associated with nystagmus is provoked by the Dix-Hallpike test. ● There is a latency period between the completion of the Dix-Hallpike test and the onset of vertigo and nystagmus. ● The provoked vertigo and nystagmus increase and then resolve within a time period of 60 seconds from onset of nystagmus.

History

Vertigo has been defined as an “illusory sensation of motion of either the self or the surroundings.” (Blakley, et al., 2001) The symptoms of vertigo resulting from posterior canal BPPV are typically described by the patient as a rotational or spinning sensation when the patient changes head position relative to gravity.

The episodes are often provoked by everyday activities and commonly occur when rolling over in bed or when the patient is tilting the head to look upward (eg, to place an object on a shelf higher than the head) or bending forward (eg, to tie shoes) (von Brevern, et al., 2007) (Furman, et al., 1999) (Dix, et al., 1952) (Whitney, et al., 2005).

Patients with BPPV most commonly report discrete, episodic periods of vertigo lasting 1 minute or less and often report modifications or limitations of their general movements to avoid provoking the vertiginous episodes (Ruckenstein, et al., 2007). Other investigators report that true “room spinning” vertigo is not always present as a reported symptom in posterior canal BPPV, with patients alternatively complaining of

lightheadedness, dizziness, nausea, or the feeling of being “off balance” (Katsarkas, et al., 1999) (von Brevern, et al., 2007) (Herdman, et al., 1997) (Herdman, et al., 1997) (Macias, et al., 2000) (Cohen, et al., 2004) (Haynes, et al., 2002) (Blatt, et al., 2000) (Norre, et al., 1995). Approximately 50 percent of patients also report subjective imbalance between the classic episodes of BPPV (von Brevern, et al., 2007). In contrast, a history of vertigo *without* associated lightheadedness may increase the a priori likelihood of a diagnosis of posterior canal BPPV (Oghalai, et al., 2000). In up to one third of cases with atypical histories of positional vertigo, Dix-Hallpike testing will still reveal positional nystagmus, strongly suggesting the diagnosis of posterior canal BPPV (Norre, et al., 1995).

Other authors have loosened the historical criteria required for BPPV diagnosis with coinage of the term “subjective BPPV” without a positive Dix-Hallpike test (Haynes, et al., 2002) (Numez, et al., 2000). However, in clinical practice, there is a practical need to balance inclusiveness of diagnosis with accuracy of diagnosis.

Physical examination

In addition to the historical criteria for the diagnosis of posterior canal BPPV, clinicians should confirm the diagnosis of posterior canal BPPV by performing the Dix-Hallpike maneuver (Table 2.1, Fig 2.1). The nystagmus produced by the Dix-Hallpike maneuvers in posterior canal BPPV typically displays two important diagnostic characteristics. First, there is a latency period between the completion of the maneuver, and the onset of subjective rotational vertigo and the objective nystagmus.

The latency period for the onset of the nystagmus with this maneuver is largely unspecified in the literature, but the panel felt that a typical latency period would range from 5 to 20 seconds, although it may be as long as 1 minute in rare cases (Baloh, et al., 1987). Second, the provoked subjective vertigo and the nystagmus increase, and then resolve within a time period of 60 seconds from the onset of nystagmus.

The fast component of the nystagmus provoked by the Dix-Hallpike maneuver demonstrates a characteristic mixed torsional and vertical movement (often described as upbeat-torsional), with the upper pole of the eye beating toward the dependent ear and the vertical component beating toward the forehead (Fig 2.1) (Furman, et al., 1999) (Honrubia, et al., 1999). Temporally, the rate of nystagmus typically begins gently, increases in intensity, and then declines in intensity as it resolves. This has been termed crescendo-decrescendo nystagmus. The nystagmus is again commonly observed after the patient returns to the upright head position and upon arising, but the direction of the nystagmus may be reversed.

Another classical feature of the nystagmus associated with posterior canal BPPV is that the nystagmus typically fatigues (a reduction in severity of nystagmus) when the maneuver is repeated (Dix, et al., 1952) (Honrubia, et al., 1999). However, repeated performance of the Dix-Hallpike maneuver to demonstrate fatigability is not recommended, because it unnecessarily subjects patients to repeated symptoms of vertigo that may be discomforting, and repeat performance may interfere with the immediate bedside treatment of BPPV (Furman, et al., 1999). Therefore, the panel did not include fatigability of the nystagmus as a diagnostic criterion.

-

Performing the Dix-Hallpike Diagnostic Maneuver

The Dix-Hallpike maneuver is performed by the clinician moving the patient through a set of specified head-positioning maneuvers to elicit the expected characteristic nystagmus of posterior canal BPPV (Fig 2.1) (Furman, et al., 1999) (Dix, et al., 1952). Before beginning the maneuver, the clinician should counsel the patient regarding the upcoming movements and warn that they may provoke a sudden onset of intense subjective vertigo,

possibly with nausea, which will subside within 60 seconds. Because the patient is going to be placed in the supine position relatively quickly with the head position slightly below the body, the patient should be oriented so that, in the supine position, the head can “hang” with support off the posterior edge of the examination. The examiner should ensure that he can support the patient’s head and guide the patient through the maneuver safely and securely, without the examiner losing support or balance himself.

1. The maneuver begins with the patient in the upright seated position with the examiner standing at the patient’s side (Furman, et al., 1999). If present, the patient’s eyeglasses should be removed. We initially describe the maneuver to test the right ear as the source of the posterior canal BPPV.
2. The examiner rotates the patient’s head 45 degrees to the right and, with manual support, maintains the 45-degree head turn to the right during the next part of the maneuver.
3. Next, the examiner fairly quickly moves the patient (who is instructed to keep the eyes open) from the seated to the supine right-ear down position and then extends the patient’s neck slightly (approximately 20 degrees below the horizontal plane) so that the patient’s chin is pointed slightly upward, with the head hanging off the edge of the examining table and supported by the examiner. The examiner observes the patient’s eyes for the latency, duration, and direction of the nystagmus (Norre, et al., 1988) (White, et al., 2005). Again, the provoked nystagmus in posterior canal BPPV is classically described as a more or less mixed torsional movement with the upper pole of both eyes beating toward the affected ear (in this example the right ear) in combination with a vertical (upbeat) component. The patient should also be queried as to the presence of subjective vertigo.
4. After resolution of the subjective vertigo and the nystagmus, if present, the patient may be slowly returned to the upright position. During the return to the upright position, a reversal of the nystagmus may be observed and should be allowed to resolve (a torsional nystagmus to the healthy ear, in combination with a vertical (downbeat) component).
5. The Dix-Hallpike maneuver (steps 1-4) should then be repeated for the left side, with the left ear arriving at the dependent position (Numez, et al., 2000). Again, the examiner should inquire about subjective vertigo and identify objective nystagmus, when present. The examination of the left side completes the test. The provoked nystagmus in left ear posterior canal BPPV is more or less mixed torsional movement with the upper pole of both eyes beating toward the affected ear (in this example the left ear) in combination with a vertical (upbeat) component. The Dix-Hallpike maneuver is considered the gold standard test for the diagnosis of posterior canal BPPV (Fife, et al., 2008). It is the most common diagnostic criterion required for entry into clinical trials and for inclusion of such trials in meta-analyses (Hilton, et al., 2004) (Cohen, et al., 2005). The lack of an alternative external gold standard to the Dix Hallpike maneuver limits the availability of rigorous sensitivity and specificity data. Although it is considered the gold standard test for posterior canal BPPV diagnosis, its accuracy may differ between specialty and nonspecialty clinicians. Lopez-Escamez et al (Lopez-Escamez, et al., 2000) have reported a sensitivity of 82 percent and specificity of 71 percent for the Dix-Hallpike maneuvers in posterior canal BPPV, primarily among specialty clinicians. In the primary care setting, Hanley and O’Dowd (Hanley, et al., 2002) have reported a positive predictive value for a positive Dix-Hallpike test of 83 percent and a negative predictive value of 52 percent for the diagnosis of BPPV. Therefore, a negative Dix-Hallpike maneuver does not necessarily rule out a diagnosis of posterior canal BPPV. Because of the lower negative predictive values of the Dix-Hallpike maneuver, it has been suggested that this maneuver may need to be repeated at a separate visit to confirm the diagnosis and avoid a false-negative result (Numez, et al., 2000) (Viire, et al., 2005) (Norre, et al., 1994). Factors that may affect the diagnostic accuracy of the Dix-Hallpike maneuver include the speed of movements during the test, time of day, and the angle of the plane of the occiput during the maneuver (Numez, et al., 2000). The Dix-Hallpike test must be done bilaterally to determine which ear is involved or if both ears are

involved (Numez, et al., 2000). In a small percent of cases, the Dix-Hallpike maneuver may be bilaterally positive (ie, the correspondingly appropriate nystagmus is elicited for each ear in the dependent position). For example, bilateral posterior canal BPPV is more likely to be encountered after head trauma (Katsarkas, et al., 1999). Although the Dix-Hallpike maneuver is the test of choice to confirm the diagnosis of posterior canal BPPV, it should be avoided in certain circumstances. Although there are no documented reports of vertebrobasilar insufficiency provoked by performing the Dix-Hallpike maneuver, clinicians should be careful to consider the risk of stroke or vascular injury in patients with significant vascular disease (Whitney, et al., 2006). Care should also be exercised in patients with cervical stenosis, severe kyphoscoliosis, limited cervical range of motion, Down syndrome, severe rheumatoid arthritis, cervical radiculopathies, Paget's disease, ankylosing spondylitis, low back dysfunction, spinal cord injuries, and morbid obesity (Whitney, et al., 2005) (Whitney, et al., 2006). Patients who are obese may be difficult for a single examiner to fully support throughout the maneuver, so additional assistance may be required. For patients with physical limitations, special tilting examination tables may allow the safe performance of the Dix-Hallpike maneuver. To our knowledge, no comparative studies have been performed so far to investigate whether the diagnostic accuracy of the Hallpike maneuver with observation of the nystagmus by the naked eye improves by the use of Frenzel's glasses or infra-red Video-Oculography.

Verantwoording

Laatst beoordeeld : 01-02-2020

Laatst geautoriseerd : 01-02-2020

Voor de volledige verantwoording, evidence tabellen en eventuele aanverwante producten raadpleegt u de Richtlijndatabase.

Referenties

- Baloh, R.W., Jacobson, K., Honrubia, V. (1993). Horizontal semicircular canal variant of benign positional vertigo. Neurology, 43, 2542-9.
- Baloh, R.W., Honrubia, V., Jacobson, K. (1987). Benign positional vertigo: clinical and oculographic features in 240 cases. Neurology, 37, 371-8.
- Blakley, B.W., Goebel, J. (2001). The meaning of the word "vertigo". Otolaryngology Head Neck Surg, 125, 147-50.
- Blatt, P.J., Georgakakis, G.A., Herdman, S.J., et al. (2000). The effect of the canalith repositioning maneuver on resolving postural instability in patients with benign paroxysmal positional vertigo. Am J Otol, 21, 356-63.
- von Brevern, M., Radtke, A., Lezius, F., et al. (2007). Epidemiology of benign paroxysmal positional vertigo: a population based study. J Neurol Neurosurg Psychiatry, 78, 710-5.
- Cakir, B.O., Cakir, B.O., Ercan, I., Cakir, Z.A., et al. (0000). What is the true incidence of horizontal semicircular canal benign paroxysmal positional vertigo. X, 0, X-.
- Califano, L., Melillo, M.G., Mazzone, S. (2010). Vassallo "Secondary signs of lateralization" in apogeotropic lateral canalolithiasis. Acta Otorhinolaryngol Ital, 30 (2), 78-86.
- Caruso, G., Nuti, D. (2005). Epidemiological data from 2270 PPV patients. Audiol Med, 3, 7-11.
- Casani, A.P., Vannucci, G., Fattori, B., et al. (2002). The treatment of horizontal canal positional vertigo: our experience in 66 cases. Laryngoscope, 112, 172-8.
- Cohen, H.S., Kimball, K.T. (2005). Effectiveness of treatments for benign paroxysmal positional vertigo of the posterior canal. Otol Neurotol, 26, 1034-40.
- Cohen, H.S. (2004). Side-lying as an alternative to the Dix-Hallpike test of the posterior canal. Otol Neurotol, 25, 130-4.
- Dix, M.R., Hallpike, C.S. (1952). The pathology, symptomatology and diagnosis of certain common disorders of the vestibular system. Ann Otol Rhinol Laryngol, 61, 987-1016.

- Eife, T.D., Iverson, D.J., Lempert, T., et al. (2008). Practice parameter: therapies for benign paroxysmal positional vertigo (an evidence-based review): Report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology*, 70, 2067-74.
- Furman, J.M., Cass, S.P. (1999). Benign paroxysmal positional vertigo. *N Engl J Med*, 341, 1590-6.
- Han, B.I., Oh, H.J., Kim, J.S. (2006). Nystagmus while recumbent in horizontal canal benign paroxysmal positional vertigo. *Neurology*, 66, 706-10.
- Hanley, K., O' Dowd, T. (2002). Symptoms of vertigo in general practice: a prospective study of diagnosis. *Br J Gen Pract*, 52, 809-12.
- Haynes, D.S., Resser, J.R., Labadie, R.F., et al. (2002). Treatment of benign positional vertigo using the semont maneuver: efficacy in patients presenting without nystagmus. *Laryngoscope*, 112, 796-801.
- Herdman, S.J. (1997). Advances in the treatment of vestibular disorders. *Phys Ther*, 77, 602-18.
- Hilton, M., Pinder, D. (2004). The Epley (canalith repositioning) manoeuvre for benign paroxysmal positional vertigo. *Cochrane Database Syst Rev*, 0, CD003162-.
- Honrubia, V., Baloh, R.W., Harris, M.R., et al. (1999). Paroxysmal positional vertigo syndrome. *Am J Otol*, 20, 465-70.
- Hornibrook, J. (2004). Horizontal canal benign positional vertigo. *Ann Otol Rhinol Laryngol*, 113, 721-5.
- Imai, T., Ito, M., Takeda, N., et al. (2005). Natural course of the remission of vertigo in patients with benign paroxysmal positional vertigo. *Neurology*, 64, 920-1.
- Katsarkas, A. (1999). Benign paroxysmal positional vertigo (BPPV): idiopathic versus post-traumatic. *Acta Otolaryngol*, 119, 745-9.
- Lopez-Escamez, J.A., Lopez-Nevot, A., Gamiz, M.J., et al. (2000). Diagnosis of common causes of vertigo using a structured clinical history. *Acta Otorrinolaringol Esp*, 51, 25-30.
- Macias, J.D., Lambert, K.M., Massingale, S., et al. (2000). Variables affecting treatment in benign paroxysmal positional vertigo. *Laryngoscope*, 110, 1921-4.
- Moon, S.Y., Kim, J.S., Kim, B.K., et al. (2006). Clinical characteristics of benign paroxysmal positional vertigo in Korea: a multicenter study. *J Korean Med Sci*, 21, 539-43.
- Norre, M.E. (1995). Reliability of examination data in the diagnosis of benign paroxysmal positional vertigo. *Am J Otol*, 16, 806-10.
- Norre, M.E. (1994). Diagnostic problems in patients with benign paroxysmal positional vertigo. *Laryngoscope*, 104, 1385-8.
- Norre, M.E., Beckers, A. (1988). Benign paroxysmal positional vertigo in the elderly. Treatment by habituation exercises.. *J Am Geriatr Soc*, 36, 425-9.
- Nunez, R.A., Cass, S.P., Furman, J.M. (2000). Short- and long-term outcomes of canalith repositioning for benign paroxysmal positional vertigo. *Otolaryngol Head Neck Surg*, 122, 647-52.
- Nuti, D., Agus, G., Barbieri, M.T., et al. (1998). The management of horizontal canal paroxysmal positional vertigo. *Acta Otolaryngol*, 118, 455-60.
- Oghalai, J.S., Manolidis, S., Barth, J.L., et al. (2000). Unrecognized benign paroxysmal positional vertigo in elderly patients. *Otolaryngol Head Neck Surg*, 122, 630-4.
- Ruckenstein, M.J., Shepard, N.T. (2007). The canalith repositioning procedure with and without mastoid oscillation for the treatment of benign paroxysmal positional vertigo. *ORL J Otorhinolaryngol Relat Spec*, 69, 295-8.
- Steenerson, R.L., Cronin, G.W., Marbach, P.M. (2005). Effectiveness of treatment techniques in 923 cases of benign paroxysmal positional vertigo. *Laryngoscope*, 115, 226-31.
- Tirelli, G., Russolo, M. (2004). 360-Degree canalith repositioning procedure for the horizontal canal. *Otolaryngol Head Neck Surg*, 131, 740-6.
- Vijire, E., Purcell, I., Baloh, R.W. (2005). The Dix-Hallpike test and the canalith repositioning maneuver. *Laryngoscope*, 115, 184-7.
- White, J., Savides, P., Cherian, N., et al. (2005). Canalith repositioning for benign paroxysmal positional vertigo. *Otol Neurotol*, 26, 704-10.
- Whitney, S.L., Morris, L.O. (2006). Multisensory impairment in older adults: evaluation and intervention. In: *Geriatric Otolaryngology*. Calhoun KH, Eibling DE, ed. New York: Taylor and Francis, 0, 115-.
- Whitney, S.L., Marchetti, G.F., Morris, L.O. (2005). Usefulness of the dizziness handicap inventory in the screening for benign paroxysmal positional vertigo. *Otol Neurotol*, 26, 1027-33.

White, J.A., Coale, K.D., Catalano, P.J., et al. (2005). Diagnosis and management of lateral semicircular canal benign paroxysmal positional vertigo. *Otolaryngol Head Neck Surg*, 133, 278-84.

De diagnose van BPPD van het horizontale kanaal

Uitgangsvraag

Wat is de beste manier om BPPD van het horizontale kanaal (h-BPPD) te diagnosticeren?

Aanbeveling

Als de patient een anamnese heeft die past bij BPPD maar de Dix-Hallpike is negatief, moet een BPPD van het horizontale kanaal worden overwogen en een supine roll test worden gedaan.

Overwegingen

- Voordeel: het onderkennen van een horizontale kanaal BPPD bij een negatieve Dix-Hallpike.
- Nadelen: het mogelijk provoceren van draaiduizeligheid
- Kosten: minimaal
- Afweging van voordeel tegen nadeel: het voordeel weegt zwaarder.
- Waarde oordelen: het belang van het onderzoeken van de patiënt op andere vormen van BPPD dan posterieur kanaal BPPD.
- Rol van de voorkeur van de patiënt: minimaal.
- Exclusie: patiënten met fysieke beperkingen




Figure 2.2

Diagrammatic views of the supine roll test. (1) The patient is in the starting neutral position. The patient's head is turned rapidly to the right side (2) to examine for characteristic nystagmus. Then the head is returned to the face-up position (1), allowing all nystagmus to subside, and then turned rapidly to the left side (3) to examine once again for nystagmus. (From Fife et al., 2008).

Onderbouwing

Conclusies

Niveau 3	De diagnose horizontale kanaal BPPD wordt gesteld wanneer draaiduizeligheid met horizontale nystagmus wordt opgewekt door de supine roll test.
-----------------	--

Samenvatting literatuur

Lateral canal BPPV (also called horizontal canal BPPV) is the second most common type of BPPV (Imai, et al., 2005) (Steenerson, et al., 2005) (Moon, et al., 2006). Because this type of BPPV has received considerably less attention in the literature, clinicians may be relatively unaware of its existence and the appropriate diagnostic maneuvers for lateral canal BPPV. Patients with a history compatible with BPPV (ie, repeated episodes of vertigo produced by changes in head position relative to gravity) who do not meet diagnostic criteria for posterior canal BPPV should be investigated for lateral canal BPPV. In many instances, the presenting symptoms of lateral canal BPPV are indistinguishable from posterior canal BPP (Streenerson, et al., 2005). Several studies have cited an incidence of approximately 10 to 15 percent in populations referred for evaluation and treatment of BPPV (White, et al., 2005) (Cakir, et al., 2006) (Hornibrook, et al., 2004) (Han, 2006) (Caruso, et al., 2005).

Furthermore, lateral canal BPPV may occur following performance of the PRMs (eg, Epley maneuver) for an initial diagnosis of posterior canal BPPV. This transition from posterior canal BPPV to lateral canal BPPV is thought to occur as free-floating particulate material migrates from the posterior canal to the lateral canal (so-called canal switch). Because this type of transition is relatively common, clinicians should be aware of lateral canal BPPV and its diagnosis (White, 2005). The supine roll test is the preferred maneuver to diagnose lateral canal BPPV (Cakir, et al., 2006) (Moon, et al., 2006) (Nutti, et al., 1998). Clinicians should inform the patient that this test is a provocative maneuver and may cause the patient to become subjectively intensely dizzy for a short period of time. The supine roll test is performed by initially positioning the patient supine with the head in neutral position followed by quickly rotating the head 90 degrees to one side with the clinician observing the patient's eyes for nystagmus (Fig 2.2). After the nystagmus subsides (or if no nystagmus is elicited), the head is then returned to the straight faceup supine position. After any additional elicited nystagmus has subsided, the head is then quickly turned 90 degrees to the opposite side, and the eyes are once again observed for nystagmus. Two potential nystagmus findings may occur with this maneuver, reflecting two types of lateral canal BPPV (White, et al., 2005) (Nutti, et al., 1998) (Tirelli, et al., 2004).

- **Geotropic type:** In most cases of lateral canal BPPV, rotation to the affected side causes a very intense horizontal nystagmus beating toward the undermost (affected) ear, known as geotropic nystagmus (ie, nystagmus with a fast component toward the ground). When the patient is rolled to the other, healthy side, there is a less intense horizontal nystagmus, again beating toward the undermost ear (again geotropic; the direction of the nystagmus has now changed).

- **Apogeotropic type:** In less common cases, rotation to the healthy side results in an intense horizontal nystagmus beating toward the uppermost (affected) ear, known as anapogeotropic nystagmus (ie, nystagmus with a fast component away from the ground). Upon rolling to the opposite (affected) side, the nystagmus will change direction, beating toward the uppermost (healthy) ear, however less intense.

In both types of lateral canal BPPV, the fast component of the strongest nystagmus always beats to the affected ear (Han, et al., 2006) (Nutti, et al., 1998) (Baloh, et al., 1993). Between the two types of lateral canal

BPPV, the geotropic variant predominates (Steenerson, et al., 2005) (Nutti, et al., 1998) (Casani, et al., 2002). Not uncommonly, because of CNS adaptation, the initially intense nystagmus may spontaneously change direction without rolling toward the opposite ear (Tirelli, et al., 2004). Also, in case of canalolithiasis, the position of the clod in the canal determines whether a geotropic (close to the utriculus) or apo-geotropic (close to the cupula) nystagmus will be induced as well, which may lead to a "spontaneous" transition from geotropic into apo-geotropic or vice-versa (Califano L, et al., 2010).

The supine roll test has not received as much widespread use or diagnostic validation as the Dix-Hallpike maneuver. Review of the literature reveals that the sensitivity and specificity of the supine roll test in the diagnosis of lateral canal BPPV have not been determined. The lack of a more accurate, commonly accepted (gold standard) test for the diagnosis of lateral canal BPPV may be responsible, in part, for the absence of data for these statistical measures. A positive supine roll test, however, is the most commonly required and consistent diagnostic entry criterion for therapeutic trials of lateral canal BPPV (Steenerson, et al., 2005) (Han, et al., 2006). Reports of harm or patient injury from the performance of the supine roll test were not identified in the literature review, although many authors simply stated that patients who could not tolerate positional maneuvers were excluded from the population under study. Care should also be exercised in patients with cervical stenosis, severe kyphoscoliosis, limited cervical range of motion, Down syndrome, severe rheumatoid arthritis, cervical radiculopathies, Paget's disease, ankylosing spondylitis, low back dysfunction, spinal cord injuries, and morbid obesity (Whitney, et al., 2005) (Whitney, et al., 2006). The benefit of performing the supine roll test is that it allows clinicians to confirm a diagnosis of lateral canal BPPV quickly and efficiently (White, et al., 2005) (Fife, et al., 2008). It also allows clinicians to more accurately and comprehensively diagnose positional vertigo that is not due to the posterior canal, whereas without supine roll testing, patients with lateral canal BPPV might be diagnostically missed if only traditional Dix-Hallpike testing was done. Further benefit might be derived from the supine roll test by decreasing the need to perform potentially unnecessary or unhelpful diagnostic testing.

Verantwoording

Laatst beoordeeld : 01-02-2020

Laatst geautoriseerd : 01-02-2020

Voor de volledige verantwoording, evidence tabellen en eventuele aanverwante producten raadpleegt u de Richtlijndatabase.

Referenties

Baloh, R.W., Jacobson, K., Honrubia, V. (1993). Horizontal semicircular canal variant of benign positional vertigo. Neurology, 43, 2542-9.

Baloh, R.W., Honrubia, V., Jacobson, K. (1987). Benign positional vertigo: clinical and oculographic features in 240 cases. Neurology, 37, 371-8.

Blakley, B.W., Goebel, J. (2001). The meaning of the word "vertigo". Otolaryngology Head Neck Surg, 125, 147-50.

Blatt, P.J., Georgakakis, G.A., Herdman, S.J., et al. (2000). The effect of the canalith repositioning maneuver on resolving postural instability in patients with benign paroxysmal positional vertigo. Am J Otol, 21, 356-63.

von Brevern, M., Radtke, A., Lezius, F., et al. (2007). Epidemiology of benign paroxysmal positional vertigo: a population based study. J Neurol Neurosurg Psychiatry, 78, 710-5.

Cakir, B.O., Ercan, I., Cakir, Z.A., et al. (2000). What is the true incidence of horizontal semicircular canal benign paroxysmal positional vertigo?. X, 0, X-

Califano, L., Melillo, M.G., Mazzone, S. (2010). Vassallo "Secondary signs of lateralization" in apogeotropic lateral

canalolithiasis. *Acta Otorhinolaryngol Ital*, 30 (2), 78-86.

Caruso, G., Nuti, D. (2005). Epidemiological data from 2270 PPV patients. *Audiol Med*, 3, 7-11.

Casani, A.P., Vannucci, G., Fattori, B., et al. (2002). The treatment of horizontal canal positional vertigo: our experience in 66 cases. *Laryngoscope*, 112, 172-8.

Cohen, H.S., Kimball, K.T. (2005). Effectiveness of treatments for benign paroxysmal positional vertigo of the posterior canal. *Otol Neurotol*, 26, 1034-40.

Cohen, H.S. (2004). Side-lying as an alternative to the Dix-Hallpike test of the posterior canal. *Otol Neurotol*, 25, 130-4.

Dix, M.R., Hallpike, C.S. (1952). The pathology, symptomatology and diagnosis of certain common disorders of the vestibular system. *Ann Otol Rhinol Laryngol*, 61, 987-1016.

Fife, T.D., Iverson, D.J., Lempert, T., et al. (2008). Practice parameter: therapies for benign paroxysmal positional vertigo (an evidence-based review): Report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology*, 70, 2067-74.

Furman, J.M., Cass, S.P. (1999). Benign paroxysmal positional vertigo. *N Engl J Med*, 341, 1590-6.

Han, B.I., Oh, H.J., Kim, J.S. (2006). Nystagmus while recumbent in horizontal canal benign paroxysmal positional vertigo. *Neurology*, 66, 706-10.

Hanley, K., O' Dowd, T. (2002). Symptoms of vertigo in general practice: a prospective study of diagnosis. *Br J Gen Pract*, 52, 809-12.

Haynes, D.S., Resser, J.R., Labadie, R.F., et al. (2002). Treatment of benign positional vertigo using the semont maneuver: efficacy in patients presenting without nystagmus. *Laryngoscope*, 112, 796-801.

Herdman, S.J. (1997). Advances in the treatment of vestibular disorders. *Phys Ther*, 77, 602-18.

Hilton, M., Pinder, D. (2004). The Epley (canalith repositioning) manoeuvre for benign paroxysmal positional vertigo. *Cochrane Database Syst Rev*, 0, CD003162-.

Honrubia, V., Baloh, R.W., Harris, M.R., et al. (1999). Paroxysmal positional vertigo syndrome. *Am J Otol*, 20, 465-70.

Hornibrook, J. (2004). Horizontal canal benign positional vertigo. *Ann Otol Rhinol Laryngol*, 113, 721-5.

Imai, T., Ito, M., Takeda, N., et al. (2005). Natural course of the remission of vertigo in patients with benign paroxysmal positional vertigo. *Neurology*, 64, 920-1.

Katsarkas, A. (1999). Benign paroxysmal positional vertigo (BPPV): idiopathic versus post-traumatic. *Acta Otolaryngol*, 119, 745-9.

Lopez-Escamez, J.A., Lopez-Nevot, A., Gamiz, M.J., et al. (2000). Diagnosis of common causes of vertigo using a structured clinical history. *Acta Otorrinolaringol Esp*, 51, 25-30.

Macias, J.D., Lambert, K.M., Massingale, S., et al. (2000). Variables affecting treatment in benign paroxysmal positional vertigo. *Laryngoscope*, 110, 1921-4.

Moon, S.Y., Kim, J.S., Kim, B.K., et al. (2006). Clinical characteristics of benign paroxysmal positional vertigo in Korea: a multicenter study. *J Korean Med Sci*, 21, 539-43.

Norre, M.E. (1995). Reliability of examination data in the diagnosis of benign paroxysmal positional vertigo. *Am J Otol*, 16, 806-10.

Norre, M.E. (1994). Diagnostic problems in patients with benign paroxysmal positional vertigo. *Laryngoscope*, 104, 1385-8.

Norre, M.E., Beckers, A. (1988). Benign paroxysmal positional vertigo in the elderly. Treatment by habituation exercises. *J Am Geriatr Soc*, 36, 425-9.

Nunez, R.A., Cass, S.P., Furman, J.M. (2000). Short- and long-term outcomes of canalith repositioning for benign paroxysmal positional vertigo. *Otolaryngol Head Neck Surg*, 122, 647-52.

Nuti, D., Agus, G., Barbieri, M.T., et al. (1998). The management of horizontal canal paroxysmal positional vertigo. *Acta Otolaryngol*, 118, 455-60.

Oghalai, J.S., Manolidis, S., Barth, J.L., et al. (2000). Unrecognized benign paroxysmal positional vertigo in elderly patients. *Otolaryngol Head Neck Surg*, 122, 630-4.

Ruckenstein, M.J., Shepard, N.T. (2007). The canalith repositioning procedure with and without mastoid oscillation for the treatment of benign paroxysmal positional vertigo. *ORL J Otorhinolaryngol Relat Spec*, 69, 295-8.

Steenerson, R.L., Cronin, G.W., Marbach, P.M. (2005). Effectiveness of treatment techniques in 923 cases of benign paroxysmal positional vertigo. *Laryngoscope*, 115, 226-31.

Tirelli, G., Russolo, M. (2004). 360-Degree canalith repositioning procedure for the horizontal canal. *Otolaryngol Head*

Neck Surg, 131, 740-6.

Virre, E., Purcell, I., Baloh, R.W. (2005). The Dix-Hallpike test and the canalith repositioning maneuver. Laryngoscope, 115, 184-7.

White, J., Savvides, P., Cherian, N., et al. (2005). Canalith repositioning for benign paroxysmal positional vertigo. Otol Neurotol, 26, 704-10.

Whitney, S.L., Morris, L.O. (2006). Multisensory impairment in older adults: evaluation and intervention. In: Geriatric Otolaryngology. Calhoun KH, Eibling DE, ed. New York: Taylor and Francis, 0, 115-.

Whitney, S.L., Marchetti, G.F., Morris, L.O. (2005). Usefulness of the dizziness handicap inventory in the screening for benign paroxysmal positional vertigo. Otol Neurotol, 26, 1027-33.

White, J.A., Coale, K.D., Catalano, P.J., et al. (2005). Diagnosis and management of lateral semicircular canal benign paroxysmal positional vertigo. Otolaryngol Head Neck Surg, 133, 278-84.

De diagnose van BPPD van het anterieure kanaal

Uitgangsvraag

Wat is de beste manier om BPPD van het anterieure kanaal (a-BPPD) te herkennen?

Aanbeveling

De diagnose a-BPPD wordt gesteld wanneer draaiduizeligheid met downbeat nystagmus wordt opgewekt door de Dix-Hallpikemanoeuvre.

Overwegingen

- Voordeel: toegenomen diagnostische accuratesse en efficiëntie.
- Nadelen: Het tijdelijk provoceren van BPPD symptomen
- Kosten: minimaal
- Afweging van voordeel tegen nadeel: de aandoening is zeer zeldzaam, zodat de procedures voor diagnostiek van anterieur kanaal BPPD niet routinematig worden uitgevoerd.
- Rol van de voorkeur van de patiënt: matig.
- Exclusie: patiënten met fysieke beperkingen zoals ernstige reumatoïde artritis en cervicale radiculopathie.

Onderbouwing

Achtergrond

Benign paroxysmal positional vertigo (BPPV) of the anterior semicircular canal (ASC) is very rare. This is probably due to the orientation of the anterior canal in the head; gravity restricts the upward movement of the statoconien debris, preventing it from entering in the canal. If debris enters the canal, gravity facilitates self-clearance through the posterior arm of the ASC into the common crus and vestibule (Korres et al., 2002). An exception is cupulolithiasis, in which debris is fixed to the cupula and can not easily leave the canal simply by gravity or 'mass inertia'.

Some studies, that did not objectivate the nystagmus with EOG or VOG, show that anterior canal types of BPPV constituted 1.6 to 12% of the cases (Celebisoy et.al., 2008, Cakir et.al., 2006, Korres et.al., 2007, Korres et.al., 2002). One study, based on EOG findings, showed a downbeat nystagmus pointing to an anterior canal BPPV in 20% of cases. This suggests that objectivating nystagmus is important for adequately diagnosing anterior canal BPPV. The optimal method to do this clinically is using infrared videorecording (Ir-VOG) that allows repeated analysis of the nystagmus type and direction without fatiguing the response by repetitive positioning manoeuvres. Anterior canal BPPV must be differentiated from central-downbeat positional nystagmus.

Conclusies

Niveau 2/3	<p>Bij patiënten met een typische anamnese voor BPPD, en een downbeat nystagmus bij de Dix Hallpike, moet de diagnose anterieur kanaal BPPD of centrale positioneringsnystagmus worden overwogen.</p> <p><i>bronnen (?)</i></p>
------------	---

Niveau 4	<p>Het is niet aangetoond, maar het lijkt waarschijnlijk dat video-oculografie behulpzaam is bij het interpreteren van de verticale component van de nystagmus.</p> <p><i>Niveau D: bronnen (niet-vergelijkend onderzoek Jackson et al 2007, en case studie Bertholon 2002))</i></p>
----------	--

Samenvatting literatuur

In the literature search we found seven case studies that described the clinical signs of anterior canal BPPV (Bertholon et.al., 2002) (Imai et.al., 2006) (Korres et.al., 2008) (Lopez-Escamez, et.al., 2006) (Ogawa et.al., 2009) (Walther et.al., 2008) (Zapala et.al., 2008). In addition we found 1 non-systematic review that described the clinical signs of BPPV (Korres et.al., 2010).

Anterior canal BPPV

The Dix-Hallpike test stimulates the contralateral anterior canal that is located in the uppermost ear during the test. In anterior canal canalolithiasis, BPPV is typically characterized by a predominantly down-beating nystagmus with a small torsional component during Dix-Hallpike testing. The possibly small torsional component may be more pronounced when the patient looks in the direction of the undermost healthy ear and is clockwise in case of a canalolithiasis of the right anterior canal and anti-clockwise in case of a canalolithiasis of the left anterior canal (Bertholon et.al., 2002, Korres et.al., 2006, 2008). Some authors report that the torsional component is often very small or absent. The direction of the torsional component of the nystagmus and the side on which vertigo and nystagmus are provoked are very important elements that may point at the affected ear.

Diagnosing the affected ear is important for treatment of the affected ear (Bertholon et al., 2002; Korres et.al., 2006, 2008).

As an alternative for the Dix-Hallpike maneuver the 'straight head-hanging' provocation maneuver is described. The head of a patient is in a middle position and the patient is placed with the head in retroflexion, stimulating both anterior canals simultaneously. Because the head is more extended during the straight head hanging position than during the Dix-Hallpike test sometimes a nystagmus may be triggered with the straight head hanging position, whereas the Dix-Hallpike test was negative. The provoked nystagmus is mostly downbeat without a clear torsional component, which makes it difficult to attribute the canalolithiasis to right or left anterior canal (Korres et al, 2008; Lopez-Escamez et al., 2006).

Tabel 2.2: criteria gebruikt voor de identificatie van het semicirculaire kanaal en de gepaste behandeling

Nystagmus/maneuver	Nystagmus duur (s)	Type	Aangedane semicirculaire kanaal
Upbeat and anti-clockwise / right Dix-Hallpike	<60	Canalolithiasis	Rechter posterior kanaal
	>60	Cupulolithiasis (Rare)	
Upbeat and clockwise / left Dix-Hallpike	<60	Canalolithiasis	Linker posterior kanaal
	>60	Cupulolithiasis (Rare)	
Downbeat and clockwise / right Dix-Hallpike	<60	Canalolithiasis	Rechter anterior kanaal
	>60	Cupulolithiasis	
Downbeat en anti-horair na linker Dix-Hallpike	<60	Canalolithiasis	Linker anterior kanaal
	>60	Cupulolithiasis	
Horizontaal geotroop/ roll test; nystagmus het sterkst na roll naar rechts.	<60	Canalolithiasis	Rechter horizontaal kanaal
	>60	Cupulolithiasis	
Horizontaal apogeotroop/ roll test; nystagmus het sterkst na roll naar rechts.	<60	Canalolithiasis	Linker horizontaal kanaal
	>60	Cupulolithiasis	

(Overgenomen uit: Anterior canal benigning paroxysmal positional vertigo, Jackson et al, 2007, die hebben het aangepast en overgenomen uit Herdman SJ Advances in the treatment of vestibular disorders. Phys Ther 1997; 77: 602-18)

Verantwoording

Laatst beoordeeld : 01-02-2020

Laatst geautoriseerd : 01-02-2020

Voor de volledige verantwoording, evidence tabellen en eventuele aanverwante producten raadpleegt u de Richtlijndatabase.

Referenties

- Bertholon, P., Bronstein, A.M., Davies, R.A., Rudge, P., Thilo, K.V. (2002). Positional down beating nystagmus in 50 patients: Cerebellar disorders and possible anterior semicircular canalolithiasis. *J Neurol Neurosurg Psychiatry*, 72, 366-372.
- Brandt, T. (2003). Benign paroxysmal positional vertigo. In: T. Brandt (ed). *Vertigo: Its Multisensory Syndromes*, 2nd edn. London: Springer, 0, 251-83.
- Cakir, B.O., Ercan, I., Cakir, Z.A., Civelek, S., Sayin, I., Turgut, S. (2006). What is the true incidence of horizontal semicircular canal benign paroxysmal positional vertigo. *Otolaryngology – Head and Neck Surgery*, 134, 451-454.
- Celebisoy, N., Polat, F., Akyurekli, O. (2008). Clinical Features of benign paroxysmal positional vertigo. *Eur Neurol*, 59, 315-319.
- Imai, T., Takeda, N., Ito, M., Nakamae, K., Sakae, H., Fujioka, H., Kubo, T. (2006). Three-dimensional analysis of benign paroxysmal positional nystagmus in a patient with anterior semicircular canal variant. *Otology & Neurotology*, 27, 362-366.
- Jackson, L.E., Morgan, B., Fletcher, J.C. Jr., Krueger, W.W. (2007). Anterior canal benign paroxysmal positional vertigo: An underappreciated entity. *Otol Neurotol*, 28 (2), 218-22.
- Korres, S., Riga, M., Sandris, V., Danielides, V., Sismanis, A. (2010). Canalolithiasis of the anterior semicircular canal (ASC): Treatment options based on the possible underlying pathogenic mechanisms. *Int J of Audiology*, 49, 606-612.
- Korres, S., Riga, M., Balatsouras, D. & Sandris, V. (2008). Benign paroxysmal positional vertigo of the anterior semicircular canal: Atypical clinical findings and possible underlying mechanisms. *Int J Audiol*, 47 (5), 276-82.
- Korres, S.G., Balatsouras, D.G., Papouliakos, S., Ferekidis, E. (2007). Benign paroxysmal positional vertigo and its management. *X*, 13 (6), CR275-282.
- Korres, S., Balatsouras, D.G., Kaberos, A., Economou, C., Kandiloros, D., et. Al. (2002). Occurrence of semicircular canal involvement in benign paroxysmal positional vertigo. *Otol. Neurology*, 23 (6), 926-32.
- Korres, S., Balatsouras, D. & Ferekidis E. (2006). Prognosis of patients with benign paroxysmal positional vertigo treated with repositioning manoeuvres. *JLO*, 120, 528-533.
- Lopez-Escamez, J., Molina, M., Gamiz, M. (2006). Anterior semicircular canal benign paroxysmal positional vertigo and positional down-beating nystagmus. *Am J Otolaryngol*, 27, 173-178.
- Ogawa, Y., Suzuki, M., Otsuka, K., Shimizu, S., Inagaki, T., Hayashi, M., Hagiwara, A., Kitajama, N. (2009). Positional and positioning down-beating nystagmus without central nervous system findings. *Auris Nasus Larynx*, 36, 698-701.
- Thalmann, R., Ignatova, E., Kachar, B., Ornitz, D.M., Thalmann, I. (2001). Development and maintenance of otoconia: biochemical considerations. *Ann N Y Acad Sci. Oct*, 942, 162-78.
- Walther, L.E., Nath, V., Krombach, G.A., Di Martino, E. (2008). Bilateral posterior semicircular canal aplasia and atypical paroxysmal positional vertigo: a case report. *Acta Otorhinolaryngologica italic*, 28, 79-82.
- Zapala, D.A. (2008). Down-beating nystagmus in anterior canal benign paroxysmal positional vertigo. *J Am Acad Audiol*, 19, 257-266.

De differentiaal diagnose van BPPD

Uitgangsvraag

Van welke andere vormen van positioneringsduizeligheid moet BPPD worden onderscheiden?

Aanbeveling

BPPD moet met name gedifferentieerd worden van andere aandoeningen die zich presenteren met houdingsafhankelijke draaiduizeligheid, zoals bijvoorbeeld orthostatische hypotensie en vestibulaire uitval.

Onderbouwing

Conclusies

Niveau 3	De differentiaaldiagnose van BPPD omvat alle aandoeningen die zich presenteren met houdingsafhankelijke duizeligheid. Hierbij moet vooral gedacht worden aan orthostatische hypotensie, centrale pathologie, angststoornissen (phobic postural vertigo) en vestibulaire migraine. Daarnaast dienen perifeer vestibulaire stoornissen, zoals bij voorbeeld M. Meniere, recurrent vestibulopathy, neuritis vestibularis en labyrinthitis, te worden overwogen.
-----------------	--

Samenvatting literatuur

Despite being the most common cause of peripheral vertigo, (Froehling, et al., 2000) BPPV is still often underdiagnosed or misdiagnosed (von Brevern, et al., 2004). Causes of vertigo that may be confused with BPPV can be divided into otological, neurological, and other entities. In subspecialty settings, BPPV caused 10.1% of vertigo, while neurological causes were rare (Bruintjes, et al., 2007).

The most common diagnoses that require distinction from BPPV are listed in Table 3.1. These conditions require distinction from BPPV because their natural history, treatment, and potential for serious medical sequelae differ significantly.

Otologic disorders

Other otological disorders causing vertigo may be differentiated from BPPV by their clinical characteristics including their temporal pattern and the presence or absence of hearing loss. Whereas BPPV is characterized by acute, discrete episodes of brief positional vertigo without associated hearing loss, other otological causes of vertigo manifest different temporal patterns and may additionally demonstrate associated hearing loss (Kentala, et al., 2003). In distinction to BPPV, Ménière's disease is characterized by discrete episodic attacks, with each attack exhibiting a characteristic triad of sustained vertigo, fluctuating hearing loss, and tinnitus (Baloh, et al., 1987), (Wladislavosky-Waserman, et al., 1984). Recurrent vestibulopathy is characterized by similar episodic attacks of vertigo, but lacks any auditory symptoms (van Leeuwen et al., 2010). As opposed to BPPV, the duration of vertigo in an episode of Ménière's disease or recurrent vestibulopathy typically lasts longer (usually on the order of hours) and is typically more disabling owing to both severity and duration. In addition, an associated contemporaneous decline in sensorineural hearing is required for the diagnosis of a Ménière's attack, whereas acute hearing loss should not occur with an episode of BPPV (Thorpe, et al., 2003). Protracted nausea and vomiting are also more common during an attack of Ménière's disease or recurrent vestibulopathy.

Acute peripheral vestibular dysfunction syndromes, such as vestibular neuritis or labyrinthitis, present with sudden, unanticipated, severe vertigo with a subjective sensation of rotational (room spinning) motion. If the auditory portion of the inner ear is affected, hearing loss and tinnitus may also result (Baloh, et al., 2003). These syndromes are commonly preceded by a viral prodrome. The time course of the vertigo is often the best differentiator between BPPV and vestibular neuritis or labyrinthitis. In vestibular neuritis or labyrinthitis, the vertigo is of gradual onset, developing over several hours, followed by a sustained level of vertigo lasting days to weeks (Kentala, et al., 2003) (Kentala, et al., 1996) (Kentala, et al., 1999). The vertigo is present at rest (not requiring positional change for its onset), but it may be subjectively exacerbated by positional changes. These acute peripheral vestibular syndromes may also be accompanied by severe levels of nausea, vomiting, sweating, and pallor, which are also typically sustained along with the vertigo.

Tabel 3.1
Differentiaaldiagnose BPPD

Otologische stoornissen	Neurologische stoornissen	Andere oorzaken
Ziekte van Ménière	Vestibulaire migraine	Angst of paniekstoornissen
Recurrent vestibulopathy Neuritis vestibularis	Vertebrobasilaire TIA's	Orthostatische hypotensie
Labyrinthitis	Demyeliniserende ziektes (MS)	Bijwerkingen van medicijnen
Superior canal dehiscence syndrome	Centraal zenuwstelsel lesies	
Posttraumatische vertigo		

Superior canal dehiscence syndrome (SCD) is clinically characterized by attacks of vertigo and oscillopsia (the sensation that viewed objects are moving or wavering back and forth) often brought on by loud sounds, Valsalva maneuvers, or pressure changes of the external auditory canals (Minor, et al., 2001). Similar to perilymphatic fistula, it differs from BPPV in that vertigo is induced by pressure changes and not position changes. SCD may also present with an associated conductive hearing loss and is diagnosed through CT of the temporal bones (Rosowski, et al., 2004).

Posttraumatic vertigo can present with a variety of clinical manifestations including vertigo, disequilibrium, tinnitus, and headache (Marzo, et al., 2004). Although BPPV is most often idiopathic, in specific cases, traumatic brain injury is associated with BPPV (Davies, et al., 1995). BPPV has been described as occurring in conjunction with or as a sequelae to other vestibular disorders as well, such as Ménière's disease and vestibular neuritis (Karlberg, et al., 2000). Therefore, clinicians must consider the possibility of more than one vestibular disorder being present in any patient who does not clearly have the specific symptoms of a single vestibular entity.

Neurological disorders

One of the key issues facing clinicians attempting to diagnose the etiology for vertigo is the differentiation between peripheral causes of vertigo (those causes arising from the ear or vestibular apparatus) and CNS causes of vertigo. Although at times this distinction may be difficult, several clinical features may suggest a central cause of vertigo rather than BPPV (Labuguen, et al., 2006) (Baloh, et al., 1998). Nystagmus findings that more strongly suggest a neurological cause for vertigo, rather than a peripheral cause such as BPPV, include

down-beating nystagmus on the Dix-Hallpike maneuver, direction-changing nystagmus occurring without changes in head position (ie, periodic alternating nystagmus), or baseline nystagmus manifesting without provocative maneuvers. Among the central causes of vertigo that should be distinguished from BPPV are migraine-associated vertigo, vertebrobasilar TIA, and intracranial tumors.

Vestibular migraine has been described as a common cause of vertigo in the adult population (Reploeg, et al., 2002) and may account for as many as 14 percent of cases of vertigo (Kentala, et al., 2003). Diagnostic criteria include 1) episodic vestibular symptoms; 2) migraine according to International Headache Society criteria; 3) at least two of the following migraine symptoms during at least two vertiginous episodes: migrainous headache, photophobia, phonophobia, or visual or other aura; and 4) other causes ruled out by appropriate investigations (Headache Classification Subcommittee of the International Headache Society, 2004). Migraine-associated vertigo is heterogeneous in that both central disorders and peripheral disorders have been described, although more often it is believed to be central in nature (Neuhauser, et al., 2001), (von Brevern, et al., 2005). It is distinguishable from BPPV by virtue of the necessary migraine/headache components, which are not associated with classic BPPV.

Several reports have suggested that isolated attacks of vertigo can be the initial and only symptom of vertebrobasilar insufficiency (Fife, et al., 1994), (Grad, et al., 1989) (Gomez, et al., 1996). Isolated transient vertigo may precede a stroke in the vertebrobasilar artery by weeks or months. The attacks of vertigo in vertebrobasilar insufficiency usually last less than 30 minutes and have no associated hearing loss. The type of nystagmus (typically gaze-evoked in central lesions), the severity of postural instability, and the presence of additional neurological signs are the main distinguishing features between vertebrobasilar insufficiency and BPPV (Gomez, et al., 1996) (Hotson, et al., 1998). In addition, the nystagmus arising in vertebrobasilar insufficiency does not fatigue and is not easily suppressed by gaze fixation, helping to separate this diagnosis from BPPV.

Intracranial tumors and other brain stem lesions may rarely present with a history and symptomatology similar to those of BPPV (Dunniway, et al., 1998). In these cases, associated symptoms such as tinnitus, aural fullness, new-onset hearing loss, and/or other neurological symptoms should help differentiate these diagnoses from BPPV. Atypical nystagmus during Dix- Hallpike testing (eg, sustained down-beating nystagmus) argues against BPPV and suggests a more serious cause. Finally, failure to respond to conservative management such as the PRM or vestibular rehabilitation should raise concern that the underlying diagnosis may not be BPPV (Dunniway, et al., 1998).

Other disorders

Several other non-otological and non-neurological disorders may present similarly to BPPV. Patients with panic disorder, anxiety disorder, or agoraphobia may complain of symptoms of lightheadedness and dizziness. Although these symptoms are usually attributed to hyperventilation, other studies have shown high prevalences of vestibular dysfunction in these patients (Jacob, et al., 1996) (Furman, et al., 2006). These conditions may also mimic BPPV. Several medications, such as Mysoline, carbamazepine, phenytoin, antihypertensive medications, and cardiovascular medications, may produce side effects of dizziness and/or vertigo and should be considered in the differential diagnosis.

Postural hypotension also may produce episodic dizziness or vertigo. The dizziness or vertigo in postural hypotension, however, is provoked by moving from the supine to the upright position in distinction to the provocative positional changes of BPPV.

Although the differential diagnosis of BPPV is vast, most of these other disorders can be further distinguished from BPPV on the basis of responses to the Dix-Hallpike maneuver and the supine roll test. Clinicians should still remain alert for concurrent diagnoses accompanying BPPV, especially in patients with a mixed clinical presentation.

Verantwoording

Laatst beoordeeld : 01-02-2020

Laatst geautoriseerd : 01-02-2020

Voor de volledige verantwoording, evidence tabellen en eventuele aanverwante producten raadpleegt u de Richtlijndatabase.

Referenties

- Baloh, R.W. (2003). Clinical practice. Vestibular neuritis. *N Engl J Med*, 348, 1027-32.
- Baloh, R.W. (1998). Differentiating between peripheral and central causes of vertigo. *Otolaryngol Head Neck Surg*, 119, 55-9.
- Baloh, R.W., Honrubia, V., Jacobson, K. (1987). Benign positional vertigo: clinical and oculographic features in 240 cases. *Neurology*, 37, 371-8.
- von Brevern, M., Zeise, D., Neuhauser, H., et al. (2005). Acute migrainous vertigo: clinical and oculographic findings. *Brain*, 128, 365-74.
- von Brevern, M., Lezius, F., Tiel-Wilck, K., et al. (2004). Benign paroxysmal positional vertigo: current status of medical management. *Otolaryngol Head Neck Surg*, 130, 381-2.
- Bruintjes, T.D., van Leeuwen, R.B. (2007). Ervaringen met multidisciplinaire benadering van duizeligheidsklachten: het Apeldoorns duizeligheidscentrum. *Ned Tijdschr KNO-Heelkunde*, 13 (4), 185-187.
- Davies, R.A., Luxon, L.M. (1995). Dizziness following head injury: a neurootological study. *J Neurol*, 242, 222-30.
- Eife, T.D., Baloh, R.W., Duckwiler, GR. (1994). Isolated dizziness in vertebrobasilar insufficiency: clinical features, angiography, and follow-up. *J Stroke Cerebrovasc Dis*, 4, 4-12.
- Dunniway, H.M., Welling, D.B. (1998). Intracranial tumors mimicking benign paroxysmal positional vertigo. *Otolaryngol Head Neck Surg*, 118, 429-36.
- Froehling, D.A., Bowen, J.M., Mohr, D.N., et al. (2000). The canalith repositioning procedure for the treatment of benign paroxysmal positional vertigo: a randomized controlled trial. *Mayo Clin Proc*, 75, 695-700.
- Furman, J.M., Redfern, M.S., Jacob, R.G. (2006). Vestibulo-ocular function in anxiety disorders. *J Vestib Res*, 16, 209-15.
- Gomez, C.R., Cruz-Flores, S., Malkoff, M.D., et al. (1996). Isolated vertigo as a manifestation of vertebrobasilar ischemia. *Neurology*, 47, 94-7.
- Grad, A., Baloh, R.W. (1989). Vertigo of vascular origin. Clinical and electronystagmographic features in 84 cases. *Arch Neurol*, 46, 281-4.
- Hanley, K., O' Dowd, T. (2002). Symptoms of vertigo in general practice: a prospective study of diagnosis. *Br J Gen Pract*, 52, 809-12.
- Headache Classification Subcommittee of the International Headache, S. (2004). The International Classification of Headache Disorders: 2nd edition. *Cephalgia*, 24, 9-160.
- Hotson, J.R., Baloh, R.W. (1998). Acute vestibular syndrome. *N Engl J Med*, 339, 680-5.
- Jacob, R.G., Furman, J.M., Durrant, J.D., et al. (1996). Panic, agoraphobia, and vestibular dysfunction. *Am J Psychiatry*, 153, 503-12.
- Karlberg, M., Hall, K., Quickert, N., et al. (2000). What inner ear diseases cause benign paroxysmal positional vertigo?. *Acta*

Otolaryngol, 120, 380-5.

Kentala, E., Rauch, S.D. (2003). A practical assessment algorithm for diagnosis of dizziness. Otolaryngol Head Neck Surg, 128, 54-9.

Kentala, E., Laurikkala, J., Pyykko, I., et al. (1999). Discovering diagnostic rules from a neurotologic database with genetic algorithms. Ann Otol Rhinol Laryngol, 108, 948-54.

Kentala, E. (1996). Characteristics of six otologic diseases involving vertigo. Am J Otol, 17, 883-92.

Van Leeuwen, R.B., Bruinjtjes, T.D. (2010). Recurrent vestibulopathy: natural course and prognostic factors. J Laryngol Otol, 124, 19-22.

Labuguen, R.H. (2006). Initial evaluation of vertigo. Am Fam Physician, 73, 244-51.

Maarsingh, O.R., Dros, J., Schellevis, F.G., van Weert, H.C., Bindels, P.J., van der Horst, HE. (2010). Dizziness reported by elderly patients in family practice: prevalence, incidence and clinical characteristics. BMC, 11, 1-9.

Marzo, S.J., Leonetti, J.P., Raffin, M.J., et al. (2004). Diagnosis and management of post-traumatic vertigo. Laryngoscope, 114, 1720-3.

Minor, L.B., Cremer, P.D., Carey, J.P., et al. (2001). Symptoms and signs in superior canal dehiscence syndrome. Ann New York Acad Sci, 942, 259-73.

Neuhauser, H., Leopold, M., von Brevern, M., et al. (2001). The interrelations of migraine, vertigo, and migrainous vertigo. Neurology, 56, 436-41.

Reploeg, M.D., , Goebel, J.A. (2002). Migraine-associated dizziness: patient characteristics and management options. Otol Neurotol, 23, 364-71.

Rosowski, J.J., Songer, J.E., Nakajima, H.H., et al. (2004). Clinical, experimental, and theoretical investigations of the effect of superior semicircular canal dehiscence on hearing mechanisms. Otol Neurotol, 25, 323-32.

Thorp, M.A., Shehab, Z.P., Bance, M.L., et al. (2003). The AAO-HNS Committee on Hearing and Equilibrium guidelines for the diagnosis and evaluation of therapy in Ménière's disease: have they been applied in the published literature of the last decade?. Clin Otolaryngol Allied Sci, 28, 173-6.

Wladislavsky-Waserman, P., Facer, G.W., Mokri, B., et al. (1984). Ménière's disease: a 30-year epidemiologic and clinical study in Rochester, MN, 1951-1980. Laryngoscope, 94, 1098-102.

Het aanvullend onderzoek van BPPD

Het onderwerp 'aanvullend onderzoek bij BPPD' wordt uitgewerkt in verschillende modules.

Verantwoording

Laatst beoordeeld : 01-02-2020

Laatst geautoriseerd : 01-02-2020

Voor de volledige verantwoording, evidence tabellen en eventuele aanverwante producten raadpleegt u de Richtlijndatabase.

Indicatie voor beeldvormend, audiologisch en vestibulair onderzoek bij BPPD

Uitgangsvraag

Wat zijn de indicaties voor beeldvormend, audiologisch en vestibulair onderzoek bij verdenking op BPPD?

Aanbeveling

Beeldvormende technieken zijn niet geïndiceerd bij de diagnose BPPD. Beeldvormende technieken dienen wel te worden toegepast bij patiënten bij wie twijfel bestaat omtrent de diagnose BPP, bijvoorbeeld als additionele neurologische uitvalssymptomen aanwezig zijn, of bij therapieresistente BPPD.

Vestibulaire functietesten hebben geen toegevoegde waarde bij patiënten met BPPD.

Vestibulaire functietesten zijn alleen geïndiceerd bij patiënten met: 1) atypische nystagmus, 2) verdenking op additionele vestibulaire pathologie 3) een falende (of herhaaldelijk falende) reactie op canalith repositiemanoeuvres (CRM), of 4) frequent recidiverende BPPD.

Overwegingen

- Voordeel: snelle behandeling mogelijk maken door het voorkomen van overbodige testen en voorkomen van mogelijke vals-positieve diagnoses; voorkomen van stralingsbelasting (MRI) en bijwerkingen door testen.
- Nadeel: potentieel missen van comorbide aandoeningen, ongemak door misselijkheid en braken ten gevolge van vestibulaire testen.
- Kosten: kostenbesparing door het voorkomen van overbodige testen.
- Afweging van voordeel tegen nadeel: het voordeel weegt zwaarder.
- Waarde oordeel: het is belangrijk om overbodige testen en vertraging in het stellen van de diagnose te voorkomen.

Onderbouwing

Conclusies

Niveau 3	Beeldvormend en vestibulair onderzoek heeft geen toegevoegde waarde bij het stellen van de diagnose BPPD.
-----------------	---

Samenvatting literatuur

The diagnosis of BPPV is based on the clinical history and physical examination. Routine radiographic imaging or vestibular testing is unnecessary in patients who already meet clinical criteria for the diagnosis of BPPV (Table 2.1). Further radiographic or vestibular testing may have a role in the diagnosis if the clinical presentation is felt to be atypical, if Dix-Hallpike testing elicits equivocal or unusual nystagmus findings, or if additional symptoms aside from those attributable to BPPV are present, suggesting an accompanying modifying CNS or otological disorder.

Radiographic Imaging

Radiographic imaging, most commonly CNS imaging using magnetic resonance or CT techniques, is commonly obtained in the evaluation of a primary symptom complaint of vertigo. However, imaging is not useful in the routine diagnosis of BPPV because there are no radiological findings characteristic of or diagnostic for BPPV (Turski, et al., 1996) (Turski, et al., 1996). The lack of characteristic findings is likely due to fact that the pathology presumed to occur in BPPV within the semicircular canals occurs at a microscopic level that is beyond the resolution of current neuroimaging techniques (Parnes, et al., 1992). On a broader scale, previous retrospective reviews of elderly patients with dizziness failed to detect any significant differences in cranial MRI findings when comparing dizzy versus non-dizzy patients (Colledge, et al., 1996) (Day, et al., 1990).

Radiographic imaging of the CNS should be reserved for patients who present with a clinical history compatible with BPPV but who also demonstrate additional neurological symptoms atypical for BPPV. Radiographic imaging may also be considered for patients with suspected BPPV but inconclusive positional testing, or in patients with other neurological signs on physical examination that are not typically associated with BPPV. Such symptoms include abnormal cranial nerve findings, visual disturbances, and severe headache, among others. It should be noted that intracranial lesions causing vertigo are rare. (Hanley, et al., 2001) Potential lesions causing vertigo identifiable on CNS imaging include cerebrovascular disease, demyelinating disease, or an intracranial mass; they are most often located in the brain stem cerebellum, thalamus, or cortex (Hanley, et al., 2001). In small case series, positional vertigo and nystagmus have been associated with neurovascular compression of cranial nerve VIII, vestibular schwannoma, Arnold Chiari malformation, and a variety of cerebellar disorders (Brandt, et al., 1994) (Jacobsen, et al., 1995) (Kumar, et al., 2002).

In distinction to standard BPPV, such conditions are quite rare and typically present with additional neurological symptoms in conjunction with the vertigo. Routine neuroimaging has not been recommended to discern these conditions from the more common causes of vertigo (Gizzi, et al., 1996). The costs of routine imaging in cases of BPPV are not justified given that diagnostic neuroimaging does not improve the diagnostic accuracy in the vast majority of BPPV cases. Therefore, neuroimaging should not be routinely used to confirm the diagnosis of BPPV.

Vestibular Function Testing

When patients meet clinical criteria for the diagnosis of BPPV (Table 2.1), no additional diagnostic benefit is obtained from vestibular function testing. Vestibular function testing is indicated when the diagnosis of a vertiginous or dizziness syndrome is unclear or possibly when the patient remains symptomatic following treatment. It may also be beneficial when multiple concurrent peripheral vestibular disorders are suspected (Baloh, et al., 1987) (Kentala, et al., 1996) (Lopez-Escamez, et al., 2003).

Vestibular function testing involves a battery of specialized tests that primarily record nystagmus in response to labyrinthine stimulation and/or voluntary eye movements. Most vestibular function testing relies on the neurological relationship between the regulation of eye movement and the balance organs: the vestibular-ocular reflex. These tests are useful in the evaluation of vestibular disorders that may not be evident from the history and clinical examination, and may provide information for quantification, prognostication, and treatment planning (Gordon, et al., 1996). The components of the vestibular function test battery identify abnormalities in ocular motility as well as deficits in labyrinthine response to position change, caloric stimulation, rotational movement, and static positions (sitting and supine). Caloric testing is an established, widely accepted technique

that is particularly useful in determining unilateral vestibular hypofunction. Rotational chair testing is considered the most sensitive and reliable technique for quantifying the magnitude of bilateral peripheral vestibular hypofunction (Fife, et al., 2008) and to assess central compensation after peripheral vestibular loss. Some or all of these test elements may be included in a vestibular test battery.

In cases of BPPV in which the nystagmus findings are suggestive but not clear, it may be beneficial to use video-oculographic recordings of nystagmus associated with posterior canal BPPV. Especially video-recorded eye movements can be analysed in detail using image-processing techniques. For further study or second opinion without the need to repeat the Dix-Hallpike manoeuvre. A second diagnostic procedure often will result in more difficult to assess eye movements because of the typical fatigue of a BPPV. In a small percentage of cases, patients with a history of positional vertigo but unclear nystagmus findings may undergo vestibular function testing. Among complex patients referred for subspecialty evaluation of BPPV, such atypical or unclear nystagmus findings may approach 13 percent in patients with diagnoses suspicious for BPPV (Bath, et al., 2000).

BPPV is relatively frequently associated with additional vestibular pathology. Symptoms associated with chronic vestibular function may persist following appropriate treatment for BPPV, even if the treatment is effective in resolving the specific complaint of positional vertigo. For example, in highly selected subsets of patients referred for subspecialty evaluation of BPPV, additional otopathology and/or vestibulopathy has been identified in 31 to 53 percent of BPPV patients (Baloh, et al., 1987) (Roberts, et al., 2005) (Korres, et al., 2004). This percentage, however, is higher than what might be expected in the nonspecialty population. Vestibular disorders that have been associated with BPPV include Ménière's disease, viral vestibular neuritis, or labyrinthitis (Hughes, et al., 1997)(Karlberg, et al., 2000). Vestibular function testing may be obtained when these additional diagnoses are suspected on the basis of signs or symptoms in addition to those of BPPV.

In patients with vestibular pathology in addition to BPPV, PRMs appear to be equally effective in resolving the positional nystagmus associated with BPPV, but complete symptom resolution is significantly less likely in those patients with additional vestibular pathology. In one study, 86 percent of patients with BPPV but without associated vestibular pathology reported complete resolution of symptoms after PRMs versus only 37 percent reporting complete resolution when additional vestibular pathology was present (Pollak, et al., 2004).

Thus, patients with suspected associated vestibular pathology in addition to BPPV may be a subset who would benefit from the additional information obtained from vestibular function testing. Similarly, up to 25 percent of patients with separate recurrences of BPPV are more likely to have associated vestibular pathology (Del Rio, et al., 2004); therefore, patients with recurrent BPPV may be candidates for vestibular function testing. In summary, patients with a clinical diagnosis of BPPV according to guideline criteria should not routinely undergo vestibular function testing, because the information provided from such testing adds little to the diagnostic accuracy in these cases, vestibular testing adds significant cost to the diagnosis and management of BPPV, and the information obtained does not alter the subsequent management of BPPV in the vast majority of the cases. Therefore, vestibular function testing should not be routinely obtained when the diagnosis of BPPV has already been confirmed by clinical diagnostic criteria. Vestibular function testing, however, may be warranted in patients with 1) atypical nystagmus, 2) suspected additional vestibular pathology, 3) a failed (or repeatedly failed) response to CRP, or 4) frequent recurrences of BPPV (Rupa, et al., 2004) (Gordon, et al., 2005).

Verantwoording

Laatst beoordeeld : 01-02-2020

Laatst geautoriseerd : 01-02-2020

Voor de volledige verantwoording, evidence tabellen en eventuele aanverwante producten raadpleegt u de Richtlijndatabase.

Referenties

- American Medical Association's Relative Value Scale Upgrade Committee (RUC) (2008). 2008 database, version 1. Based on 2005-06 Medicare Part B data; Centers for Medicare and Medicaid Services. Chicago: American Medical Association, 0, X-.
- Baloh, R.W., Honrubia, V., Jacobson, K. (1987). Benign positional vertigo: clinical and oculographic features in 240 cases. *Neurology*, 37, 371-8.
- Bath, A.P., Walsh, R.M., Ranalli, P., et al. (2000). Experience from a multidisciplinary "dizzy" clinic. *Am J Otol*, 21, 92-7.
- Brandt, T., Dieterich, M. (1994). VIIIth nerve vascular compression syndrome: vestibular paroxysmia. *Baillieres Clin Neurol*, 3, 565-75.
- Colledge, N.R., Barr-Hamilton, R.M., Lewis, S.J., et al. (1996). Evaluation of investigations to diagnose the cause of dizziness in elderly people: a community based controlled study. *BMJ*, 313, 788-92.
- Day, J.J., Freer, C.E., Dixon, A.K., et al. (1990). Magnetic resonance imaging of the brain and brain-stem in elderly patients with dizziness. *Age Ageing*, 19, 144-50.
- Eife, T.D., Tusa, R.J., Furman, J.M., et al. (2000). Assessment: vestibular testing techniques in adults and children: report of the Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology. *Neurology*, 55, 1431-41.
- Jacobson, G., Butcher, J.A., Newman, C.W., et al. (1995). When paroxysmal positional vertigo isn't benign. *J Am Acad Audiol*, 6, 346-9.
- Gizzi, M., Riley, E., Molinari, S. (1996). The diagnostic value of imaging the patient with dizziness. A Bayesian approach. *Arch Neurol*, 53, 1299-304.
- Gordon, C.R., Shupak, A., Spitzer, O., et al. (1996). Nonspecific vertigo with normal otoneurological examination. The role of vestibular laboratory tests. *J Laryngol Otol*, 110, 1133-7.
- Gordon, C.R., Gadoth, N. (2005). Benign paroxysmal positional vertigo: who can diagnose it, how should it be treated and where. *Harefuah*, 144, 567-71.
- Hanley, K., O'Dowd, T., Considine, N. (2001). A systematic review of vertigo in primary care. *Br J Gen Pract*, 51, 666-71.
- Havlik, R.J. (1986). Aging in the eighties, impaired senses for sound and light in persons aged 65 years and over, preliminary data from the supplement on aging to the national health interview survey, United States, January-June 1984. *Advanced Data*. *Vital Health Stat*, 125, 2-.
- Hughes, C.A., Proctor, L. (1997). Benign paroxysmal positional vertigo. *Laryngoscope*, 107, 607-13.
- Karlberg, M., Hall, K., Quickert, N., et al. (2000). What inner ear diseases cause benign paroxysmal positional vertigo?. *Acta Otolaryngol*, 120, 380-5.
- Kentala, E., Rauch, S.D. (2003). A practical assessment algorithm for diagnosis of dizziness. *Otolaryngol Head Neck Surg*, 128, 54-9.
- Kentala, E., Viikki, K., Pyykko, I., et al. (2000). Production of diagnostic rules from a neurotologic database with decision trees. *Ann Otol Rhinol Laryngol*, 109, 170-6.
- Kentala, E., Pyykko, I. (2000). Vertigo in patients with benign paroxysmal positional vertigo. *Acta Otolaryngol*, 543, 20-2.
- Kentala, E., Laurikkala, J., Pyykko, I., et al. (1999). Discovering diagnostic rules from a neurotologic database with genetic algorithms. *Ann Otol Rhinol Laryngol*, 108, 948-54.
- Kentala, E. (1996). Characteristics of six otologic diseases involving vertigo. *Am J Otol*, 17, 883-92.
- Korres, S.G., Balatsouras, D.G. (2004). Diagnostic, pathophysiologic, and therapeutic aspects of benign paroxysmal positional vertigo. *Otolaryngol Head Neck Surg*, 131, 438-44.
- Kumar, A., Patni, A.H., Charbel, F. (2002). The Chiari I malformation and the neurotologist. *Otol Neurotol*, 23, 727-35.
- Parnes, L.S., Agrawal, S.K., Atlas, J. (2003). Diagnosis and management of benign paroxysmal positional vertigo (BPPV). *CMAJ*, 169, 681-93.
- Pollak, L., Davies, R.A., Luxon, L.L. (2002). Effectiveness of the particle repositioning maneuver in benign paroxysmal positional vertigo with and without additional vestibular pathology. *Otol Neurotol*, 23, 79-83.

- Del Rio, M., Arriaga, M.A. (2004). Benign positional vertigo: prognostic factors. *Otolaryngol Head Neck Surg*, 130, 426-9.
- Roberts, R.A., Gans, R.E., Kastner, A.H., et al. (2005). Prevalence of vestibulopathy in benign paroxysmal positional vertigo patients with and without prior otologic history. *Int J Audiol*, 44, 191-6.
- Rupa, V. (2004). Persistent vertigo following particle repositioning maneuvers: an analysis of causes. *Arch Otolaryngol Head Neck Surg*, 130, 436-9.
- Stewart, M.G., Chen, A.Y., Wyatt, J.R., et al. (1999). Cost-effectiveness of the diagnostic evaluation of vertigo. *Laryngoscope*, 109, 600-5.
- Turski, P., Seidenwurm, D., Davis, P., et al. (1996). American College of Radiology: ACR appropriateness criteria: vertigo and hearing loss. Reston (VA): American College of Radiology, 0, 8-.
- Turski, P., Seidenwurm, D., Davis, P., et al. (2006). American College of Radiology: Expert Panel on Neuroimaging: vertigo and hearing loss. Reston (VA): American College of Radiology, 0, 8-.

Indicatie voor audiometrisch onderzoek bij BPPD

Uitgangsvraag

Kunnen audiometrische testen de diagnose BPPD ondersteunen?

Aanbeveling

Er is bij BPPD geen indicatie voor het uitvoeren van audiometrisch onderzoek.

Overwegingen

- Voordeel: geen vertraging in herkenning en behandeling van BPPD
- Nadeel: mogelijk missen van gehoorverlies (passend bij bijv. M. Meniere)
- Kosten: mogelijke realisatie van kostenbesparingen als er minder audiogrammen worden aangevraagd.
- Afweging van voordeel tegen nadeel: het voordeel weegt zwaarder.
- Waarde oordeel: Het is gemakkelijk om een kleine groep patiënten te identificeren waarbij audiometrie nuttig zou zijn op basis van de anamnese.
- Rol van de voorkeur van de patiënt: minimaal.

Onderbouwing

Conclusies

Niveau 4	Audiometrie heeft geen toegevoegde waarde om BPPD te diagnosticeren.
-----------------	--

Samenvatting literatuur

Audiometry is the most commonly obtained objective test of hearing. Recent Medicare data indicate that approximately 9 percent of audiograms obtained annually are ordered in association with diagnostic categories related to vertigo (International Classification of Diseases, Version 9 codes: 386 and/or 780.4) (American Medical association's relative value scale upgrade, 2008). Specialty clinicians with access to audiometry frequently obtain audiometry as part of the evaluation of vertigo in contradistinction to nonspecialty clinicians. However, limited diagnostic cohort studies and cost-effectiveness studies supporting this practice are available.

Audiometry is not required to diagnose BPPV; however, audiometry may offer some diagnostic benefit for patients in whom the clinical diagnosis of BPPV is unclear. Both hearing loss and BPPV are more prevalent in older patients. Therefore, BPPV and some degree of hearing loss (likely long-standing, as in presbycusis) are likely to coexist in patients with BPPV (Havlik, et al., 1986). From a pathophysiological standpoint, a preexisting, stable hearing loss should be unrelated to and not influence the diagnosis of BPPV. In such cases, routine audiometry is unlikely to reinforce or influence the diagnosis of BPPV.

In the majority of cohort studies of BPPV, audiometric studies, when obtained, have been largely normal. In some of these studies, however, the inclusion criteria for a diagnosis of BPPV included no history of antecedent hearing loss (Kentala, et al., 2000). In two algorithmic studies, audiometry was found to be cost-effective and

diagnostically effective in the broad evaluation of patients with vertigo (Kentala, et al., 2000) (Kentala, et al., 2003). In a study of 192 patients referred to an academic center for the evaluation of vertigo, Stewart et al (Stewart, et al., 1999) found that the audiogram was the most cost-effective test among various studies including electronystagmography, posturography, MRI, and blood tests. Notably, however, the cost-effectiveness (diagnostic benefit) of the history and physical examination (ie, Dix- Hallpike maneuver or supine role test) was not directly studied. This diagnostic focus notably differs from the current guideline, which emphasizes the value of the clinical history and physical examination.

In a study of 564 cases, Kentala et al (Kentala, et al., 1999) found in a diagnostic algorithm analysis that the presence of a normal audiogram was corroborating for a diagnosis of BPPV, distinguishing BPPV from other associated conditions such as Ménière's disease, vestibular schwannoma, and so on. However, the panel felt that distinction from such associated conditions could be made accurately and more cost-effectively on the basis of the history, rather than relying on audiometry. Upon review of the literature, no meaningful observational or diagnostic cohort studies either supporting or arguing against the use of audiometry in the diagnosis of the BPPV population was identified.

Traditional BPPV should not manifest with symptoms of a new-onset hearing loss. A newly reported hearing loss arising in conjunction with vertigo suggests a diagnosis other than BPPV and such patients merit audiometry. Clinicians should distinguish patients with vertigo and newonset hearing loss from those patients with preexisting otological disease who subsequently develop BPPV. As noted, studies have reported rates of associated otological or vestibular pathology in 30 to 50 percent of cases in referred populations with BPPV (Baloh, et al., 1987) (Roberts, et al., 2005) (Korres, et al., 2004). In cases with preexisting otological disease and a diagnostic concern for BPPV, audiometry may help establish the independent stability of the otological disease, thereby helping to confirm a diagnosis of BPPV.

Audiometry is a noninvasive test with widespread availability and no reported harms from testing. The potential benefits of obtaining audiometry in the evaluation of BPPV include the ability to establish baseline stability or, alternatively, to help rule out other otological conditions such as Ménière's disease or labyrinthitis (Kentala, et al., 1999). The primary disadvantage of routinely obtaining audiometry in patients undergoing evaluation for BPPV is clearly the cost to the health care system. In the vast majority of cases of BPPV with stable hearing by history, the audiogram is most likely to be normal or demonstrate an age-appropriate sensorineural hearing loss and, therefore, likely will not influence the diagnosis of BPPV. Overall, insufficient evidence exists to either confirm or disaffirm the value of routine audiometry in the initial assessment of BPPV.

Verantwoording

Laatst beoordeeld : 01-02-2020

Laatst geautoriseerd : 01-02-2020

Voor de volledige verantwoording, evidence tabellen en eventuele aanverwante producten raadpleegt u de Richtlijndatabase.

Referenties

American Medical Association's Relative Value Scale Upgrade Committee (RUC) (2008). 2008 database, version 1. Based

on 2005-06 Medicare Part B data; Centers for Medicare and Medicaid Services. Chicago: American Medical Association, 0, X-.

Baloh, R.W., Honrubia, V., Jacobson, K. (1987). Benign positional vertigo: clinical and oculographic features in 240 cases. *Neurology*, 37, 371-8.

Bath, A.P., Walsh, R.M., Ranalli, P., et al. (2000). Experience from a multidisciplinary “dizzy” clinic. *Am J Otol*, 21, 92-7.

Brandt, T., Dieterich, M. (1994). VIIIth nerve vascular compression syndrome: vestibular paroxysmia. *Baillieres Clin Neurol*, 3, 565-75.

Colledge, N.R., Barr-Hamilton, R.M., Lewis, S.J., et al. (1996). Evaluation of investigations to diagnose the cause of dizziness in elderly people: a community based controlled study. *BMJ*, 313, 788-92.

Day, J.J., Freer, C.E., Dixon, A.K., et al. (1990). Magnetic resonance imaging of the brain and brain-stem in elderly patients with dizziness. *Age Ageing*, 19, 144-50.

Eife, T.D., Tusa, R.J., Furman, J.M., et al. (2000). Assessment: vestibular testing techniques in adults and children: report of the Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology. *Neurology*, 55, 1431-41.

Jacobson, G., Butcher, J.A., Newman, C.W., et al. (1995). When paroxysmal positional vertigo isn't benign. *J Am Acad Audiol*, 6, 346-9.

Gizzi, M., Riley, E., Molinari, S. (1996). The diagnostic value of imaging the patient with dizziness. A Bayesian approach. *Arch Neurol*, 53, 1299-304.

Gordon, C.R., Shupak, A., Spitzer, O., et al. (1996). Nonspecific vertigo with normal otoneurological examination. The role of vestibular laboratory tests. *J Laryngol Otol*, 110, 1133-7.

Gordon, C.R., Gadoth, N. (2005). Benign paroxysmal positional vertigo: who can diagnose it, how should it be treated and where. *Harefuah*, 144, 567-71.

Hanley, K., O'Dowd, T., Considine, N. (2001). A systematic review of vertigo in primary care. *Br J Gen Pract*, 51, 666-71.

Havlik, R.J. (1986). Aging in the eighties, impaired senses for sound and light in persons aged 65 years and over, preliminary data from the supplement on aging to the national health interview survey, United States, January-June 1984. *Advanced Data. . Vital Health Stat*, 125, 2-.

Hughes, C.A., Proctor, L. (1997). Benign paroxysmal positional vertigo. *Laryngoscope*, 107, 607-13.

Karlberg, M., Hall, K., Quickert, N., et al. (2000). What inner ear diseases cause benign paroxysmal positional vertigo?. *Acta Otolaryngol*, 120, 380-5.

Kentala, E., Rauch, S.D. (2003). A practical assessment algorithm for diagnosis of dizziness. *Otolaryngol Head Neck Surg*, 128, 54-9.

Kentala, E., Viikki, K., Pyykko, I., et al. (2000). Production of diagnostic rules from a neurotologic database with decision trees. *Ann Otol Rhinol Laryngol*, 109, 170-6.

Kentala, E., Pyykko, I. (2000). Vertigo in patients with benign paroxysmal positional vertigo. *Acta Otolaryngol*, 543, 20-2.

Kentala, E., Laurikkala, J., Pyykko, I., et al. (1999). Discovering diagnostic rules from a neurotologic database with genetic algorithms. *Ann Otol Rhinol Laryngol*, 108, 948-54.

Kentala, E. (1996). Characteristics of six otologic diseases involving vertigo. *Am J Otol*, 17, 883-92.

Korres, S.G., Balatsouras, D.G. (2004). Diagnostic, pathophysiologic, and therapeutic aspects of benign paroxysmal positional vertigo. *Otolaryngol Head Neck Surg*, 131, 438-44.

Kumar, A., Patni, A.H., Charbel, F. (2002). The Chiari I malformation and the neurotologist. *Otol Neurotol*, 23, 727-35.

Parnes, L.S., Agrawal, S.K., Atlas, J. (2003). Diagnosis and management of benign paroxysmal positional vertigo (BPPV). *CMAJ*, 169, 681-93.

Pollak, L., Davies, R.A., Luxon, L.L. (2002). Effectiveness of the particle repositioning maneuver in benign paroxysmal positional vertigo with and without additional vestibular pathology. *Otol Neurotol*, 23, 79-83.

Del Rio, M., Arriaga, M.A. (2004). Benign positional vertigo: prognostic factors. *Otolaryngol Head Neck Surg*, 130, 426-9.

Roberts, R.A., Gans, R.E., Kastner, A.H., et al. (2005). Prevalence of vestibulopathy in benign paroxysmal positional vertigo patients with and without prior otologic history. *Int J Audiol*, 44, 191-6.

Rupa, V. (2004). Persistent vertigo following particle repositioning maneuvers: an analysis of causes. *Arch Otolaryngol Head Neck Surg*, 130, 436-9.

Stewart, M.G., Chen, A.Y., Wyatt, J.R., et al. (1999). Cost-effectiveness of the diagnostic evaluation of vertigo.

Laryngoscope, 109, 600-5.

Turski, P., Seidenwurm, D., Davis, P., et al. (1996). American College of Radiology: ACR appropriateness criteria: vertigo and hearing loss. Reston (VA): American College of Radiology, 0, 8-.

Turski, P., Seidenwurm, D., Davis, P., et al. (2006). American College of Radiology: Expert Panel on Neuroimaging: vertigo and hearing loss. Reston (VA): American College of Radiology, 0, 8-.

De behandeling van BPPD

Het onderwerp 'behandeling van BPPD' wordt uitgewerkt in verschillende modules.

Verantwoording

Laatst beoordeeld : 01-02-2020

Laatst geautoriseerd : 01-02-2020

Voor de volledige verantwoording, evidence tabellen en eventuele aanverwante producten raadpleegt u de Richtlijndatabase.

Repositiemanoeuvres als behandeling van BPPD

Uitgangsvraag

Zijn repositiemanoeuvres geschikt als therapie om patiënten met BPPD te behandelen?

Aanbeveling

BPPD posterieure kanaal

Behandel objectieve en subjectieve BPPD van het posterieure kanaal met een Epley- of een Semontmanoeuvre.

Behandel objectieve en subjectieve BPPD van het posterieure kanaal door een arts of specifiek daartoe geschoold persoon, zoals een fysiotherapeut of klinisch fysisch laborant.

Raad de patiënt in principe niet aan om de Epley- of Semontmanoeuvre zelf uit te voeren.

BPPD horizontale kanaal

Behandel patiënten met BPPD van het horizontale kanaal (geotrope variant) met een Gufonimaneuvre of de barbecuemanoeuvre (Lempert).

Verwijs patiënten met verdenking op BPPD van het horizontale kanaal (apogeotrope variant) naar een centrum of collega met veel ervaring in de behandeling van BPPD.

Behandel patiënten met BPPD van het horizontale kanaal door een arts of specifiek daartoe geschoold persoon, zoals een fysiotherapeut of klinisch fysisch laborant.

BPPD anterieure kanaal

Verwijs patiënten met verdenking op BPPD van het anterieure kanaal naar een centrum of collega met veel ervaring in de behandeling van BPPD.

Overwegingen

Voor de diagnose van BPPD worden de internationale criteria aangehouden die zijn geformuleerd door de 'Committee for Classification of Vestibular Disorders of the Bárány Society' (Von Brevern et al., 2015).

Posterieur kanaal BPPD

Er kan onderscheid gemaakt worden tussen verschillende typen p-BPPD (zie tabel 1). De diagnose p-BPPD is zeker als bij de Dix-Hallpike test zowel duizeligheid als nystagmus kan worden opgewekt. Er wordt dan gesproken van 'objectieve BPPD'. Als er bij de Dix-Hallpike wel duizeligheid, maar geen nystagmus kan worden opgewekt, wordt de term 'subjectieve BPPD' gebruikt. De behandeling van beide types BPPD is hetzelfde. Als de anamnese typisch is voor BPPD, maar bij de Dix-Hallpike test geen duizeligheid of nystagmus kan worden opgewekt, is herbeoordeling op een ander tijdstip (met herhaling van de Dix-Hallpike test) gewenst (Alvarenga et al., 2011; Balatsouras et al., 2012; Haynes et al., 2002; Huebner et al., 2013; Jung et al., 2016; Tirelli et al., 2001; Weider et al., 1994).

Tabel 1: onderscheid tussen de verschillende typen p-BPPD.

	Anamnestic verdacht	Dix Hallpike test	
		Nystagmus	Duizeligheid
Objectieve p-BPPD (definitive)	+	+	+
Subjectieve p-BPPD (possible)	+	-	+
Anamnestic p-BPPD (probable)	+	-	-

Behandeling van posterieur kanaal BPPD geschiedt bij voorkeur met een Epley- of Semontmanoeuvre. De effectiviteit van deze manoeuvres is het meeste onderzocht. Daarnaast is al jarenlang ervaring met deze manoeuvres opgedaan, de kans op complicaties is uitermate klein en de kwaliteit van leven lijkt ook te verbeteren (Gupta, 2018). De belangrijkste complicatie is conversie naar de horizontale kanaal BPPD-variant. Beide manoeuvres zijn makkelijk en snel uit te voeren en brengen weinig kosten met zich mee. De manoeuvres zijn uitvoerbaar door de behandelend arts of een daartoe geschoold persoon (bijvoorbeeld fysiotherapeut, KNF-laborant).

In het algemeen wordt het thuis, door de patiënt zelf, uitvoeren van een Epleymanoeuvre niet aanbevolen. De patiënt moet namelijk precies weten welke kant behandeld dient te worden en hoe de manoeuvre uitgevoerd moet worden. Het is ook nog niet aangetoond dat een 'thuis-Epley' even effectief is als een Epley uitgevoerd door een zorgprofessional. Daarnaast is de toegang tot hulpverleners, bijvoorbeeld fysiotherapeuten, in Nederland makkelijk. Patiënten kunnen er snel naartoe, zonder verwijzing. In uitzonderlijke gevallen en op basis van klinische expertise kan een 'thuis-Epley' wel zinvol zijn, bijvoorbeeld bij patiënten die al lang bekend zijn met het ziektebeeld, de zijdigheid van de aandoening kennen en getraind zijn in het uitvoeren van de Epley.

In de behandeling van BPPD zijn er ook andere manoeuvres beschikbaar, zoals de Quick Liberatory Rotation en Li Quick manoeuvre, maar deze manoeuvres zijn nog niet zo bekend en de bewijskracht voor deze behandelmethodes is minder groot dan voor de Epley en Semont.

Horizontaal kanaal BPPD

Geotrope variant

Behandeling van de geotrope variant van horizontaal kanaal BPPD geschiedt bij voorkeur met een Gufoni- of barbecuemanooeuvre (Lempert). Deze behandelingen zijn het meest bekend en hebben de hoogste bewijskracht voor wat betreft effectiviteit. Ze zijn makkelijk uitvoerbaar en gaan niet gepaard met hoge kosten. De behandeling geeft weinig complicaties.

De 'Li Quick repositiemanoeuvre' is mogelijk ook effectief in de behandeling van de geotrope variant van h-BPPD, maar deze verdient niet de voorkeur aangezien deze behandeling minder goed is onderzocht en de effectiviteit nog niet duidelijk is aangetoond.

Apogeotrope variant

De apogeotrope variant van horizontaal kanaal BPPD komt minder vaak voor dan de geotrope variant en de behandeling is minder goed onderzocht. Het advies is daarom bij verdenking op de apogeotrope variant van h-BPPD door te verwijzen naar een centrum of collega met veel ervaring in de behandeling van BPPD.

Complicerende factor is namelijk dat men vooraf niet weet waar het otoconiaal debris zich precies in het semicirculaire kanaal bevindt en of het debris los zit in het kanaal (canalolithiasis) of vastzit aan de cupula (cupulolithiasis). In de behandeling van de apogeotrope variant lijkt een 'head-shaking' manoeuvre effectiever dan de Semontmanoeuvre. De gemodificeerde Gufonimanoeuvere en mastoidoscillatie lijken effectiever dan een shammanoeuvre in het verdwijnen van vertigo en nystagmus. Echter voor het goed uitvoeren van deze manoeuvres is uitgebreidere expertise nodig.

Anterieur kanaal BPPD

Anterieur kanaal BPPD is uitermate zeldzaam. Behandeling van a-BPPD is gebaseerd op expert opinion en observationeel onderzoek. Het advies is om patiënten met een verdenking op a-BPPD daarom door te sturen naar een centrum of collega met veel ervaring in de behandeling van BPPD.

Onderbouwing

Achtergrond

Canalith repositiemanoeuvres kunnen zorgen voor een aanzienlijke vermindering van de duizeligheidssymptomen bij benigne paroxysmale positie duizeligheid (BPPD). De effectiviteit van de Epley- en Semontmanoeuvre bij BPPD van het posterieure semicirculaire kanaal (p-BPPD) is bekend. Sinds 2011 zijn er nieuwe repositiemanoeuvres beschikbaar, zowel voor BPPD van het posterieure als het horizontale kanaal en is er meer bekend over de behandeling van BPPD van het horizontale kanaal (h-BPPD). Specifiek voor h-BPPD, wordt onderscheid gemaakt tussen de geotrope en apogeotrope vorm (afhankelijk van de richting van de nystagmus tijdens de zogenaamde 'supine roll test') en hangt de behandeling af van het gevonden type. In deze module wordt uitgezocht welke repositiemanoeuvres geschikt zijn als therapie voor patiënten met BPPD.

Conclusies

Posterieur kanaal BPPD

Uitkomstmaat: verdwijnen van symptomen

<p>Redelijk GRADE</p>	<p>In de behandeling van BPPD van het posterieure kanaal zijn de Epley- en Semontmanoeuvre effectief in het verdwijnen van de symptomen.</p> <p><i>Bronnen: (Zhang, 2017; Hilton, 2014; van Duijn, 2014; Saberi, 2017; Li, 2017; Mandalà, 2012; Karanjai, 2010)</i></p>
<p>Laag GRADE</p>	<p>In de behandeling van BPPD van het posterieure kanaal is mogelijk de Li Quick repositiemanoeuvre effectief in het verdwijnen van de symptomen.</p> <p><i>Bronnen: (Li, 2017)xxx)</i></p>

Uitkomstmaat: conversie naar een negatieve Dix-Hallpike

Redelijk GRADE	<p>In de behandeling van BPPD van het posterieure kanaal zijn de Epley- en de Semontmanoeuvre effectief in de conversie naar een negatieve uitkomst in de Dix-Hallpike test.</p> <p><i>Bronnen: (Liu, 2016; Hilton, 2014; Van Duijn, 2014; Helminski, 2010; Saberi, 2017; Califano, 2003)</i></p>
---------------------------	---

Laag GRADE	<p>In de behandeling van BPPD van het posterieure kanaal is mogelijk de Quick Liberatory Rotation effectief in de conversie naar een negatieve uitkomst in de Dix-Hallpike test.</p> <p><i>Bronnen: (Califano, 2003)</i></p>
-----------------------	--

Horizontaal kanaal BPPD - geotrope variant

Uitkomstmaat: verdwijnen van symptomen

Laag GRADE	<p>In de behandeling van de geotrope variant van BPPD van het horizontale kanaal zijn de Gufonimanoevrre, de barbecuemanoeuvre en de 'Li Quick repositiemanoeuvre' effectief in het verdwijnen van de symptomen.</p> <p><i>Bronnen: (Li, 2018; Testa, 2012; Van den Broek, 2014)</i></p>
-----------------------	--

Horizontaal kanaal BPPD - apogeotrope variant

Uitkomstmaat: verdwijnen van symptomen

Laag GRADE	<p>In de behandeling van de apogeotrope variant lijkt een 'head-shaking' manoeuvre effectiever dan de Semontmanoeuvre. De gemodificeerde Gufonimanoevrre en mastoidoscillatie lijken effectiever dan een shammanoeuvre in het verdwijnen van de symptomen.</p> <p><i>Bronnen: (Kim, 2017; Kim, 2012a; Oh, 2009)</i></p>
-----------------------	---

Anterieur kanaal BPPD

- GRADE	Geen vergelijkend onderzoek heeft de behandeling van anterieur kanaal BPPD bekeken.
--------------------	---

Uitkomstmaat: kwaliteit van leven

- GRADE	Geen enkele studie heeft kwaliteit van leven meegenomen als uitkomstmaat, daarom kunnen daar geen conclusies over worden geformuleerd.
--------------------	--

Samenvatting literatuur

Voor het bestuderen van de effectiviteit van repositiemanoeuvres als behandeling van Benigne Paroxysmale Positie Duizeligheid (BPPD) is het ideale studiedesign een RCT waarin de effectiviteit van een repositiemanoeuvre zoals de Epleymanoeuvre, de barbecuemanoeuvre of de Gufonimanoevrre bij patiënten met BPPD wordt vergeleken met geen behandeling, placebo of een andere repositiemanoeuvre. Allereerst zijn

er zes systematische reviews geïnccludeerd met betrekking tot de probleemstelling (op basis van RCT's). Daarnaast zijn er negen RCT's geïnccludeerd die nog niet zijn opgenomen in de geïnccludeerde systematische reviews. De meeste studies rapporteerden verdwijnen van symptomen en/of een conversie naar een negatieve Dix-Hallpike als uitkomstmaten. Geen enkele studie heeft kwaliteit van leven meegenomen als uitkomstmaat.

De studies waren heterogeen wat betreft het type interventie en het subtype BPPD waarbij de interventie werd ingezet; om deze reden was er geen pooling van data mogelijk. De resultaten van de studies zijn daarom afzonderlijk beschreven.

Beschrijving studies

Systematic reviews

Six systematic reviews were selected based on full text.

PC-BPPV

Four SRs performed a systematic review in which the efficacy of the Epley manoeuvre was compared with the Semont's Liberatory manoeuvre (Zhang, 2017; Liu, 2016) or placebo (Hilton, 2014) or watchful waiting (van Duijn, 2014) in patients with PC-BPPV.

One SR compared the canalith repositioning procedure (CRP) with the Liberatory manoeuvre or sham in patients with PC-BPPV (Helminski, 2010).

HC-BPPV (geotropic)

One SR described results of RCTs comparing the Gufoni manoeuvre compared to the barbecue or sham manoeuvre in patients with the geotropic type of HC-BPPV (van den Broek, 2014).

Outcome measures were recovery rate (Zhang, 2017), a negative Dix-Hallpike test (Liu, 2016; Hilton 2014; van Duijn, 2014; Helminski, 2010), complete resolution of vertigo symptoms (Hilton, 2014), adverse effects of treatment (Hilton, 2014) and success percentage, defined as the resolution of vertigo (van den Broek, 2014).

RCTs

Posterior canal BPPV (PC-BPPV)

Saberi (2017) performed a randomized clinical trial in patients with PC-BPPV. Patients were randomized over two groups; 30 patients received the Epley manoeuvre and were compared to another group of 30 patients receiving the Gans manoeuvre.

Li (2017) performed a single blinded prospective controlled study on 120 patients with PC-BPPV. Patients were randomly assigned to the Li quick repositioning manoeuvre or the Epley manoeuvre.

Mandalà (2012) performed a double-blind randomized trial on the short-term efficacy of the Semont's Liberatory Manoeuvre (SLM) for the treatment of PC-BPPV. A total of 342 patients with unilateral PC-BPPV were recruited for a multicenter study. Patients were randomly assigned to treatment by SLM (n = 174) or sham treatment (n = 168). Subjects were followed up twice (1 and 24 h) with the Dix-Hallpike manoeuvre by blinded examiners.

Karanjai (2010) performed a prospective randomised clinical trial in which they compared the Semont's manoeuvre, the Epley manoeuvre and Brandt-Daroff exercises in the treatment of BPPV. In total 48 patients diagnosed with PC-BPPV were randomized over the three different treatment groups. Follow-up was at 2 weeks and 3 months.

Califano (2003) performed a randomized trial to compare three treatment options for BPPV. Three hundred patients with posterior canal/cupulolithias were divided into 3 treatment groups: 100 treated by Semont Technique; 100 by a Repositioning procedure (Parnes technique); 100 by a new manoeuvre called "Quick Liberatory Rotation".

Horizontal canal BPPV (HC-BPPV) (geotropic)

Li (2018) performed a single blinded prospective controlled study on 120 patients with geotropic HC-BPPV. Patients were randomly assigned to the Li quick repositioning manoeuvre (n=60) or the barbecue repositioning manoeuvre (n=60).

Testa (2012) performed a randomized controlled trial to compare the effectiveness of a new technical variant of the Gufoni manoeuvre as treatment of HC-BPPV. Patients were randomized over the two different treatment groups (N=87).

Horizontal canal BPPV (HC-BPPV) (apogeotropic)

Kim (2017) performed a randomized, prospective, sham-controlled study to determine the immediate and short-term effect of mastoid oscillation versus modified Gufoni manoeuvre in patients with the apogeotropic type of HC-BPPV. Patients were randomly assigned to receive a single application of Gufoni (n= 70), mastoid oscillation (n= 67), or sham manoeuvre (n= 72).

Kim (2012a) performed a randomized, prospective, sham-controlled study to determine the immediate and long-term therapeutic efficacies of Gufoni and head-shaking manoeuvres in apogeotropic type of HC-BPPV. Patients were randomized to Gufoni (n= 52), head-shaking (n=54), or sham maneuver (n= 51).

Oh (2009) performed a prospective randomized trial to compare therapeutic head-shaking and the modified Semont manoeuvre in apogeotropic HC-BPPV. Patients were alternately assigned to each treatment arm (N=103).

Anterior canal BPPV (AC-BPPV)

There were no studies that described manoeuvres for the treatment of AC-BPPV.

Resultaten

SRs

PC-BPPV

Outcome: resolution of symptoms

Zhang (2017) concluded that the Semont's liberatory manoeuvre is as effective as the Epley Manoeuvre and Brandt-Daroff Exercises for PC-BPPV treatment.

Outcome: conversion to negative Dix-Hallpike

Liu (2016) concluded that the Epley manoeuvre was similar to the Semont manoeuvre in both efficacy and safety for PC-BPPV in short-term effects, and both were superior to the sham treatment. However, in the long run the Epley manoeuvre being superior to the Semont manoeuvre (based upon five included studies).

Helminski (2010) concluded that randomized controlled trials provided evidence that the Canalith Repositioning Procedure (CRP) resolves PC-BPPV, and quasi-RCTs suggested that the CRP or the Liberatory Manoeuvre performed by a clinician or with proper instruction at home by the patient resolves PC-BPPV. There were no data on the effects of the manoeuvres on outcomes relevant to patients.

Both outcomes reported

Van Duijn (2014) concluded that all data of the selected studies show a benefit in favour of the Epley manoeuvre at 1-week follow-up in the management of PC-BPPV.

Conclusions made in the Cochrane review by Hilton (2014) concluded that there is evidence that the Epley manoeuvre is a safe, effective treatment for PC- BPPV, based on the results of 11, mostly small, RCTs with relatively short follow-up. The recurrence rate of BPPV after initial successful treatment with the modified Epley manoeuvre is 36%, although evidence is conflicting and based on only two small trials. Outcomes for Epley manoeuvre treatment are comparable to treatment with Semont and Gans manoeuvres, but superior to Brandt-Daroff exercises.

HC-BPPV (geotropic)

Outcome: resolution of symptoms

Van de Broek (2014) concluded that given the variety in the treatments compared and follow-up duration, all three studies included showed that the Gufoni manoeuvre was more effective than the sham manoeuvre or treatment with vestibular suppressants in patients with HC-BPPV. All three studies mentioned that the manoeuvre was easy to perform, which made it suitable for older, immobile, and obese patients. However, there were insufficient data to establish the relative efficacy of the Gufoni manoeuvre compared with other manoeuvres.

RCTs

PC-BPPV

Outcome: resolution of symptoms

Saberi (2017) reported that after one day of intervention by Epley manoeuvre, 86.7% of patients (n= 26) showed subjective improvement, whereas 60% of patients (n= 18) treated by Gans manoeuvre reported subjective response; the difference was statistically significant ($p= 0.02$).

Li (2017) reported that there was no statistically significant difference between the Li manoeuvre and the Epley manoeuvre group in terms of the 3-day and 1-week success rates of treatment. In the Li manoeuvre group (n=56), 35 (62.5%) and 49 (87.5%) exhibited successful repositioning at 3 days (one repositioning cycle) and 1 week (two repositioning cycles) after the first repositioning, respectively. Seven (12.5%) failed to respond to two

cycles treatment. In the Epley manoeuvre group ($n=57$), 34 (59.6%) and 52 cases (91.2%) exhibited successful repositioning at 3 days and 1 week, respectively, after the first repositioning. Five (8.8%) failed to respond to two cycles treatment.

Mandalà (2012) showed that at the 1 and 24 h follow-up, 138 and 151 (79.3 and 86.8%), respectively, patients undergoing the Semont's Liberatory Manoeuvre (SLM) had recovered from vertigo, compared to none of the patients undergoing the sham manoeuvre ($p<0.0001$).

Karanjai (2010) showed that at two weeks, the reported resolution of symptoms was 75% ($n=12$) in the Semont's group, 87.6% ($n=14$) in the Epley group and 56.25% ($n=9$) in the group who underwent Brandt-Daroff exercises. The authors conclude that performance of any of the three manoeuvres can be expected to give good results in the management of BPPV and that Epley manoeuvre appears to give better results than the other two. However, no statistical analysis was performed to confirm this conclusion.

Outcome: conversion to negative Dix-Hallpike

Saberi (2017) showed that one day after the intervention a significant objective improvement was found among Epley-treated group compared with the Gans group (86.7%, $n=26$ versus 56.7%, $n=17$, $p=0.01$). Bivariate analysis indicated that the relative chances of subjective and objective responses to Epley manoeuvre rather than Gans after 1 day were 4.32 (95% CI 1.2 to 15.6) and 4.97 (95% CI 1.38 to 17.8) and after 1 week were 2.66 folds (95% CI 0.92 to 7.69). For the long term this study revealed similar efficacy of Epley and Gans manoeuvre for the treatment of BPPV. Cervical pain was most frequent complication of Epley manoeuvre.

Califano (2003) showed that success rates of the different treatments were 97% ($n=97$) for the Semont's manoeuvre, 98% ($n=98$) for the Quick Liberatory Manoeuvre and 96% ($n=96$) for the Canalith Repositioning Manoeuvre. The primary outcome was the absence of paroxysmal positional nystagmus induced by Dix-Hallpike. There were no significant differences between the groups.

HC-BPPV (geotropic)

Outcome: resolution of symptoms

Li (2018) reported successful repositioning rates of 53.3% ($n=32$), 70.4% ($n=38$), 90.7% ($n=49$) and 92.3% ($n=48$) in barbecue manoeuvre group and 61.7% ($n=37$), 80.7% ($n=46$), 93.0% ($n=53$) and 96.3% ($n=52$), in Li manoeuvre group at 1-day, 3-day, 1-week and 1-month follow-up respectively. Differences in success rates of repositioning between Li and barbecue manoeuvre groups at 1 day, 3 days, 1 week and 1 month after initial treatment were not statistically significant using the Kaplan–Meier survival curve with a log-rank test ($p=0.270$). This study shows that there are no significant differences in short-term efficacy between the Li quick positioning manoeuvre and the classic barbecue manoeuvre in the treatment of HC-BPPV. However, as the Li manoeuvre is rapid, well-tolerated and does not require specific body positions, it can be used for treating patients who are elderly, obese, intolerant of slow manoeuvres, or who have a limited range of motion of the cervical spine.

Testa (2012) reported that after the modified Gufoni, 41 patients (93%) were cured after the first session of which 40 patients (91%) had a complete resolution of symptoms without conversion to PC-BPPV and 1 patient (2%) had a conversion to PC-BPPV and was successfully treated with Semont's manoeuvre. Three patients (7%) did not show any benefit after the treatment. After treatment with Gufoni 38 patients (88%) were cured of which

31 patients (72%) had a complete resolution of vertigo and nystagmus without conversion to PC-BPPV and 7 patients (16%) had a conversion to PC-BPPV, and were successfully treated with Semont's manoeuvre. Five patients (12%) did not have any benefit after the treatment. The percentage of symptoms resolution after the first session with the modified Gufoni's manoeuvre was not statistically different than those observed with the Gufoni's manoeuvre (93% versus 88%), but the modified Gufoni's manoeuvre appears more effective than Gufoni's manoeuvre ($\chi^2 = 6.13$, $P = 0.047$) to reduce the percentage of conversion of HSC-BPPV into PSC-BPPV (2% versus 16%).

HC-BPPV - apogeotropic

Outcome: resolution of symptoms

Kim (2017) reported that the immediate responses of the modified Gufoni manoeuvre (33/70, 47.1%) and mastoid oscillation (32/67, 47.8%) were better than the sham manoeuvre (14/72, 19.4%) ($p < 0.0001$). The second-day results did not differ among the three groups ($p = 0.76$). The short-term responses showed better efficacies with the Gufoni manoeuvre (51/70, 76.1%) and mastoid oscillation (46/67, 71.9%) than with the sham manoeuvre (38/72, 53.5%) ($p = 0.02$). Therapeutic efficacies did not differ between the Gufoni and mastoid oscillation groups in terms of both immediate and short-term outcomes ($p = 0.94$, 0.57).

Kim (2012a) showed that after a maximum of two manoeuvres on the initial visit day, Gufoni (38/52, 73.1%) and head-shaking (33/53, 62.3%) manoeuvres showed better responses than the sham maneuver (17/49, 34.7%). The cumulative therapeutic effects were also better with Gufoni ($p < 0.001$) and head-shaking ($p = 0.026$) manoeuvres compared with the sham maneuver. However, therapeutic efficacies did not differ between the Gufoni and head-shaking groups in terms of both immediate ($p = 0.129$) and long-term ($p = 0.239$) outcomes.

Oh (2009) showed that in the head-shaking group 37.7% of the patients showed resolution of vertigo and nystagmus ($n=17$) or transition of apogeotropic into geotropic HC-BPPV ($n=2$). In the modified Semont group only 17.3% of the patients showed resolution of vertigo and nystagmus ($n=7$) or transition into the geotropic HC-BPPV. So, the head-shaking manoeuvre described by the authors proved more effective than the modified Semont manoeuvre in treating apogeotropic HC-BPPV.

Overzicht resultaten per subtype BPPD*			
		<i>Gemaakte vergelijking</i>	<i>Resultaten en conclusies</i>
P-BPPD			
Zhang (2017)	SR	Semont versus. Epley en/of versus. Brandt-Daroff	Semont en Epley even effectief en effectiever dan Brandt-Daroff
Liu (2016)	SR	Semont versus. Epley	Epley en Semont even effectief en effectiever dan Sham
Hilton (2014)	SR	Epley versus. placebo	Epley is effectief
Van Duijn (2014)	SR	Epley versus afwachtend beleid	Epley is effectief

Helminski (2010)	SR	Canalith repositiemanoeuvre versus. Sham/Liberatory	CRP en LM zijn beide effectief
Saberi (2017)	RCT	Epley versus. Gans	Op korte termijn is Epley effectiever dan Gans, op lange termijn even effectief
Li (2017)	RCT	Li quick repositiemanoeuvre versus. Epley	Li en Epley beide effectief, geen verschil tussen beide behandelingen
Mandalà (2012)	RCT	Semont versus. Sham	Semont is effectief
Karanjai (2010)	RCT	Semont versus. Epley versus. Brandt-Daroff	Semont, Epley en Brandt-Daroff zijn effectief. Epley lijkt effectiever dan de andere twee behandelingen, maar statistische onderbouwing ontbreekt
Califano (2003)	RCT	Semont versus. Canalith repositiemanoeuvre (Parnes) versus. Quick Liberatory Rotation	Semont, Canalith repositiemanoeuvre en Quick alledrie effectief

H-BPPD - geotrope variant

Li (2018)	RCT	Li quick repositiemanoeuvre versus. barbecue	Beide effectief, geen verschil
Testa (2012)**	RCT	Modified Gufoni versus. Gufoni	Beide effectief, geen verschil
Van den Broek (2014)	SR	Gufoni versus. barbecue/sham	Gufoni effectiever dan sham. Onduidelijk of Gufoni ook beter is dan barbecue

H-BPPD - apogeotrope variant

Kim (2017)	RCT	Modified Gufoni versus mastoid oscillation versus. sham	Modified Gufoni en mastoid oscillation effectiever dan sham, geen verschil tussen modified Gufoni en mastoid oscillation
Kim (2012a)	RCT	Gufoni versus. Head-shaking versus. sham	Gufoni en head-shaking effectiever dan sham, geen verschil tussen Gufoni en head-shaking
Oh (2009)	RCT	Therapeutisch 'head-shaking' versus. Semont	'Head-shaking' effectiever dan Semont

*in de studies worden verschillende termen voor dezelfde manoeuvres gebruikt. De Canalith repositiemanoeuvre is hetzelfde als de Epleymanoeuvre en de Semontmanoeuvre is hetzelfde als de Liberatory manoeuvre.

**subtype niet gespecificeerd naar geotrope/apogeotrope variant

Bewijskracht van de literatuur

De kracht van het wetenschappelijk bewijs werd bepaald volgens de GRADE-methode en is gerapporteerd per uitkomstmaat.

Posterieur kanaal BPPD

Epley- en Semontmanoeuvre

De bewijskracht voor de uitkomstmaten 'verdwijnen van symptomen' en 'conversie naar een negatieve Dix-Hallpike' is met 1 niveau verlaagd van hoog naar redelijk bewijs gezien de beperkingen in heterogeniteit van de geïnccludeerde studies (inconsistentie).

Li Quick repositiemaneuvre en Quick Liberatory Rotation,

De bewijskracht voor de uitkomstmaten 'verdwijnen van symptomen' en 'conversie naar een negatieve Dix-Hallpike' is met 2 niveaus verlaagd van hoog naar laag bewijs gezien beperkingen in de onderzoeksopzet (risk of bias) en het geringe aantal patiënten (imprecisie).

Horizontaal kanaal BPPD

Gufonimaneuvre, gemodificeerde Gufonimaneuvre, barbecuemanoeuvre, Li Quick repositiemaneuvre, head-shaking, mastoidoscillatie

De bewijskracht voor de uitkomstmaten 'verdwijnen van symptomen', is met 2 niveaus verlaagd van hoog naar laag bewijs gezien beperkingen in de onderzoeksopzet (risk of bias) en heterogeniteit van de geïnccludeerde studies (inconsistentie).

Zoeken en selecteren

Om de uitgangsvraag te kunnen beantwoorden is er een systematische literatuuranalyse verricht naar de volgende zoekvraag:

Wat is de effectiviteit van repositiemaneuvres, zoals de Epleymanoeuvre, barbecuemanoeuvre (Lempert) of de Gufonimaneuvre bij patiënten met BPPD in vergelijking met geen behandeling of placebo?

P: patiënten met BPPD, verschillende groepen:

1. posterieur kanaal BPPD;
- 2a. horizontaal kanaal BPPD, geotrope variant;
- 2b. horizontaal kanaal BPPD, apogeotrope variant;
3. anterieur kanaal BPPD.

I:

1. Epleymanoeuvre;
- 2a. barbecuemanoeuvre (Lempert), Gufonimaneuvre;
- 2b. gemodificeerde Gufonimaneuvre, head-shakemanoeuvre;
3. reversed Epley manoeuvre, Yacovinomaneuvre.

C:

1. geen behandeling/placebo/Brandt-Daroff/Semontmanoeuvre (bevrijdingsmanoeuvre)/Foster manoeuvre/Somersault manoeuvre;
2. geen behandeling/placebo/andere manoeuvres;
3. geen behandeling/placebo/andere manoeuvres.

O: verdwijnen van symptomen, kwaliteit van leven, conversie naar een negatieve Dix-Hallpike.

Relevante uitkomstmaten

De werkgroep achtte verdwijnen van symptomen, kwaliteit van leven en conversie Dix-Hallpike voor de besluitvorming cruciale uitkomstmaten.

De werkgroep definieerde niet a priori de genoemde uitkomstmaten, maar hanteerde de in de studies gebruikte definities.

Zoeken en selecteren (Methode)

In de databases Medline (via OVID) en Embase is op 7 februari 2019 met relevante zoektermen gezocht naar systematische reviews (SRs) en randomized controlled trials (RCT's). De zoekverantwoording is weergegeven onder het tabblad Verantwoording. De literatuurzoekactie leverde 300 treffers op (63 SR en 237 RCT's). Studies werden geselecteerd op grond van de volgende selectiecriteria: vergelijkende studies waarin de effectiviteit van repositiemanoeuvres bij BPPD wordt vergeleken met placebo of met elkaar. Op basis van titel en abstract werden in eerste instantie 67 studies voorgeselecteerd (17 SRs en 50 RCT's). Na raadpleging van de volledige tekst, werden vervolgens 51 studies geëxcludeerd (11 SRs en 40 RCT's) (zie exclusietabel onder het tabblad Verantwoording) en 16 studies definitief geselecteerd (6 SRs en 10 RCT's).

Resultaten

Zes SRs en tien RCT's zijn opgenomen in de literatuuranalyse. De belangrijkste studiekarakteristieken en resultaten zijn opgenomen in de evidencetabellen. De beoordeling van de individuele studieopzet (risk of bias (RoB)) is opgenomen in de risk-of-biastabellen.

Verantwoording

Laatst beoordeeld : 01-02-2020

Laatst geautoriseerd : 01-02-2020

Voor de volledige verantwoording, evidence tabellen en eventuele aanverwante producten raadpleegt u de Richtlijndatabase.

Referenties

- Alvarenga, G. A., Barbosa, M. A., & Porto, C. C. (2011). Benign paroxysmal positional vertigo without nystagmus: diagnosis and treatment. *Brazilian journal of otorhinolaryngology*, 77(6), 799-804.
- Balatsouras, D. G., & Korres, S. G. (2012). Subjective benign paroxysmal positional vertigo. *Otolaryngology--Head and Neck Surgery*, 146(1), 98-103.
- Califano, L., Capparuccia, P. G. G., Di Maria, D., Melillo, M. G., & Villari, D. (2003). Treatment of benign paroxysmal positional vertigo of posterior semicircular canal by " Quick Liberatory Rotation Manoeuvre ". *Acta otorhinolaryngologica italica*, 23(3), 161-167.
- Fife, T.D., Iverson, D.J., Lempert, T., Furman, J.M., Baloh, R.W., Tusa, R.J., Hain, T.C., Herdman, S., Morrow, M.J., Gronseth, G.S. (2008) Practice parameter: therapies for benign positional vertigo (an evidence-based review). Report of the quality standards subcommittee of the American academy of neurology. *Neurology*; 70, 2067-2074.
- Gupta, A. K., Sharma, K. G., & Sharma, P. (2019). Effect of Epley, Semont Maneuvers and Brandt–Daroff Exercise on Quality of Life in Patients with Posterior Semicircular Canal Benign Paroxysmal Positional Vertigo (PSCBPPV). *Indian Journal of Otolaryngology and Head & Neck Surgery*, 71(1), 99-103.

- Haynes, D. S., Resser, J. R., Labadie, R. F., Girasole, C. R., Kovach, B. T., Scheker, L. E., & Walker, D. C. (2002). Treatment of benign positional vertigo using the semont maneuver: efficacy in patients presenting without nystagmus. *The Laryngoscope*, 112(5), 796-801.
- Helminski, J. O., Zee, D. S., Janssen, I., & Hain, T. C. (2010). Effectiveness of particle repositioning manoeuvres in the treatment of benign paroxysmal positional vertigo: a systematic review. *Physical therapy*, 90(5), 663-678.
- Hilton, M. P., & Pinder, D. K. (2014). The Epley (canalith repositioning) manoeuvre for benign paroxysmal positional vertigo. *Cochrane database of systematic reviews*, (12).
- Huebner, A. C., Lytle, S. R., Doettl, S. M., Plyler, P. N., & Thelin, J. T. (2013). Treatment of objective and subjective benign paroxysmal positional vertigo. *Journal of the American Academy of Audiology*, 24(7), 600-606.
- Jung, J. Y., & Kim, S. H. (2016). Comparison between objective and subjective benign paroxysmal positional vertigo: clinical features and outcomes. *Acta oto-laryngologica*, 136(12), 1267-1272.
- Karanjai, S., & Saha, A. K. (2010). Evaluation of vestibular exercises in the management of benign paroxysmal positional vertigo. *Indian Journal of Otolaryngology and Head & Neck Surgery*, 62(2), 202-207.
- Kim, J. S., Oh, S. Y., Lee, S. H., Kang, J. H., Kim, D. U., Jeong, S. H., ... & Kim, H. J. (2012). Randomized clinical trial for apogeotropic horizontal canal benign paroxysmal positional vertigo. *Neurology*, 78(3), 159-166.
- Kim, H. A., Park, S. W., Kim, J., Kang, B. G., Lee, J., Han, B. I., ... & Lee, H. (2017). Efficacy of mastoid oscillation and the Gufoni maneuver for treating apogeotropic horizontal benign positional vertigo: a randomized controlled study. *Journal of neurology*, 264(5), 848-855.
- Li, J., Zou, S., & Tian, S. (2018). A prospective randomized controlled study of Li quick repositioning manoeuvre for geotropic horizontal canal BPPV. *Acta oto-laryngologica*, 138(9), 779-784.
- Li, J., Tian, S., & Zou, S. (2017). Efficacy of the Li manoeuvre in treating posterior canal benign paroxysmal positional vertigo. *Acta oto-laryngologica*, 137(6), 588-592.
- Liu, Y., Wang, W., Zhang, A. B., Bai, X., & Zhang, S. (2016). Epley and Semont manoeuvres for posterior canal benign paroxysmal positional vertigo: A network meta-analysis. *The Laryngoscope*, 126(4), 951-955.
- Mandalà, M., Santoro, G. P., Libonati, G. A., Casani, A. P., Faralli, M., Giannoni, B., ... & Vannucchi, P. (2012). Double-blind randomized trial on short-term efficacy of the Semont manoeuvre for the treatment of posterior canal benign paroxysmal positional vertigo. *Journal of neurology*, 259(5), 882-885.
- Oh, S. Y., Kim, J. S., Jeong, S. H., Oh, Y. M., Choi, K. D., Kim, B. K., ... & Lee, J. J. (2009). Treatment of apogeotropic benign positional vertigo: comparison of therapeutic head-shaking and modified Semont manoeuvre. *Journal of neurology*, 256(8), 1330-1336.
- Saberi, A., Nemati, S., Sabnan, S., Mollahoseini, F., & Kazemnejad, E. (2017). A safe-repositioning manoeuvre for the management of benign paroxysmal positional vertigo: Gans versus Epley manoeuvre; a randomized comparative clinical trial. *European Archives of Oto-Rhino-Laryngology*, 274(8), 2973-2979.
- Testa, D., Castaldo, G., De Santis, C., Trusio, A., & Motta, G. (2012). Treatment of horizontal canal benign paroxysmal positional vertigo: a new rehabilitation technique. *The Scientific World Journal*, 2012.
- Tirelli, G., D'Orlando, E., Giacomarra, V., & Russolo, M. (2001). Benign positional vertigo without detectable nystagmus. *The Laryngoscope*, 111(6), 1053-1056.
- van den Broek, E. M. J. M., van der Zaag-Loonen, H. J., & Brintjes, T. D. (2014). Systematic review: efficacy of gufoni manoeuvre for treatment of lateral canal benign paroxysmal positional vertigo with geotropic nystagmus. *Otolaryngology--Head and Neck Surgery*, 150(6), 933-938.
- van Duijn, J. G., Isfordink, L. M., Nij Bijvank, J. A., Stapper, C. W., van Vuren, A. J., Wegner, I., ... & Grolman, W. (2014). Rapid systematic review of the epley manoeuvre for treating posterior canal benign paroxysmal positional vertigo. *Otolaryngology--Head and Neck Surgery*, 150(6), 925-932.
- Von Brevern, M., Bertholon, P., Brandt, T., Fife, T., Imai, T., Nuti, D., & Newman-Toker, D. (2015). Benign paroxysmal positional vertigo: diagnostic criteria. *Journal of Vestibular Research*, 25(3, 4), 105-117.
- Weider, D. J., Ryder, C. J., & Stram, J. R. (1994). Benign paroxysmal positional vertigo: analysis of 44 cases treated by the canalith repositioning procedure of Epley. *The American journal of otology*, 15(3), 321-326.
- Zhang, X., Qian, X., Lu, L., Chen, J., Liu, J., Lin, C., & Gao, X. (2017). Effects of Semont manoeuvre on benign paroxysmal positional vertigo: a meta-analysis. *Acta oto-laryngologica*, 137(1), 63-70.

Vestibulaire revalidatie als behandeling van BPPD

Uitgangsvraag

Is vestibulaire revalidatie geschikt als therapie om patiënten met BPPD te behandelen?

Aanbeveling

Vestibulaire revalidatie is niet primair geïndiceerd bij BPPD.

Overwegingen

- Voordeel: potentieel sneller verlost van symptomen dan met observatie alleen.
- Nadeel: geen ernstige bijwerkingen gezien in de gepubliceerde trials, provocatie van BPPD symptomen door revalidatie oefeningen, mogelijk een tragere oplossing dan repositie manoeuvres.
- Kosten: noodzaak tot herhaalde bezoeken
- Afwegingen: de voordelen wegen niet op tegen de nadelen, wel kan bij ouderen vestibulaire revalidatie worden overwogen, als aanvullende therapie na een repositiemanoeuvre.
- Waarde oordeel: vestibulaire revalidatie is mogelijk beter geschikt als aanvullende therapie dan als primaire behandeling (een subgroep van patiënten met balansstoornissen, centraal zenuwstelsel stoornissen of risico om te vallen, kunnen meer profiteren van vestibulaire revalidatie dan patiënten met uitsluitend BPPD).
- Rol van de voorkeur van de patiënt: aanzienlijk gezien de gezamenlijke beslissing
- Exclusiecriteria: patiënten met fysieke beperkingen

Onderbouwing

Conclusies

Niveau 2/3	<p>Vestibulaire revalidatie in de vorm van adaptatieoefeningen is bij posterieure kanaal BPPD effectief in vergelijking met placebo. Op korte termijn lijkt vestibulaire revalidatie minder effectief dan een repositiemanoeuvre, maar op lange termijn is vestibulaire revalidatie mogelijk net zo effectief als een repositiemanoeuvre. Er zijn aanwijzingen dat, vooral bij ouderen, vestibulaire revalidatie als aanvullende therapie naast een repositiemanoeuvre beter zou beschermen tegen het terugkeren van de klachten dan een repositiemanoeuvre alleen.</p> <p>Er zijn onvoldoende gegevens om de effectiviteit van vestibulaire therapie bij horizontale kanaal BPPD te beoordelen.</p>
-------------------	--

Samenvatting literatuur

The clinician may offer vestibular rehabilitation, either self-administered or with a clinician, for the initial treatment of BPPV. Option based on controlled observational studies and a balance of benefit and harm. Overview of Vestibular Therapy Vestibular rehabilitation is a form of physical therapy designed to promote habituation, adaptation, and compensation for deficits related to a wide variety of balance disorders. It may also be referred to as vestibular habituation, vestibular exercises, or vestibular therapy. There is no single specific protocol for vestibular rehabilitation, but rather a program of therapy is developed on the basis of the underlying diagnosis. Programs can include canalith repositioning exercises, adaptation exercises for gaze

stabilization, habituation exercises, substitution training for visual or somatosensory input, postural control exercises, fall prevention training, relaxation training, conditioning exercises, functional skills retraining, and patient and family education (Herdman, et al., 2000) (Telian, et al., 1996) (Whitney, et al., 2000).

With respect to BPPV, vestibular rehabilitation programs most commonly focus on habituation exercises either in formal outpatient therapy programs or with home exercise programs. Vestibular rehabilitation programs may also include PRMs, but repositioning maneuvers will be covered separately in the guideline. Herein, we refer to vestibular rehabilitation as a series of exercises or training maneuvers performed by the patient for the treatment of BPPV with or without direct clinician supervision.

Vestibular rehabilitation habituation exercises were first described by Cawthorne and Cooksey in the 1940s (Cawthorne, et al., 1944). These exercises consist of a series of eye, head, and body movements in a hierarchy of increasing difficulty, which provokes vestibular symptoms. The exercises begin with simple head movements, performed in the sitting or supine position, and progress to complex activities, including walking on slopes and steps with eyes open and closed, and sports activities requiring eye-hand coordination. These exercises theoretically fatigue the vestibular response and force the CNS to compensate by habituation to the stimulus (Norre, et al., 1987a/b). In 1980, Brandt and Daroff (Brandt, et al., 1980) (Brandt, et al., 1994) described home repositioning exercises that involve a sequence of rapid lateral head/trunk tilts repeated serially to promote loosening and ultimately dispersion of debris toward the utricular cavity. In these exercises, the patient starts in a sitting position and moves quickly to the right-side lying position, with the head rotated 45 degrees and facing upward. This position is maintained for 30 seconds after the vertigo stops. The patient then moves rapidly to a left-side lying position, with the head rotated 45 degrees and facing upward. In early work with patients with BPPV, patients repeated these maneuvers moving from the sitting to side-lying position three times a day for 2 weeks while hospitalized and had excellent resolution of BPPV symptoms (Troost, et al., 1992).

Vestibular Rehabilitation as a Treatment of BPPV

Relatively few RCTs and case series have been published regarding the effectiveness of vestibular rehabilitation as the initial therapy for BPPV. In a prospective analysis of 25 consecutive patients with BPPV, Banfield et al (Banfield, et al., 2000) reported that patients demonstrate an excellent short-term response rate of 96 percent subjectively to vestibular rehabilitation treatment with an average of three clinic visits per patient, but the authors noted a significant recurrence rate of BPPV with long-term follow-up (mean follow-up 3.8 years). The authors cited one advantage of vestibular rehabilitation: the capability of patients to be self-reliant in their ability to return to habituation exercises should symptoms recur. In a controlled trial of 60 patients with BPPV comparing a PRM, vestibular rehabilitation exercises and no treatment, vestibular rehabilitation provided better resolution of vertigo compared with no treatment (Steenerson, et al., 1996). The PRM arm demonstrated resolution of symptoms with fewer treatments than those required for vestibular rehabilitation, although the relative improvements at 3-month follow-up were comparable.

Several studies have compared vestibular rehabilitation exercises to particle rehabilitation maneuvers in the treatment of posterior canal BPPV. In an RCT of 124 patients randomized to CRP, modified liberatory maneuver, sham maneuver, Brandt-Daroff exercises, and vestibular habituation exercises by Cohen, repositioning maneuvers were more effective than Brandt-Daroff exercises or habituation exercises (Cohen et.al., 2005). Both types of vestibular rehabilitation treatments, however, were individually more effective than a

sham intervention (Cohen, et al., 2005)(Hillier, 2007). Soto Varela et al (Soto Varela, et al., 2001) comparatively analyzed a total of 106 BPPV patients randomly assigned to receive Brandt-Daroff habituation exercises, the Semont maneuver, or the Epley maneuver. At the 1-week follow-up, similar cure rates were obtained with the Semont and Epley maneuvers (74% and 71%, respectively), both cure rates being significantly higher than that obtained with Brandt- Daroff exercises (24%). At 3-month follow-up, the cure rate for the Brandt-Daroff exercises increased significantly to 62 percent, although the rate was still lower than that of PRMs. Other studies have demonstrated similar results for vestibular rehabilitation in BPPV (Furman, 1999) (Toledo, 2000). In a double blind RCT control study Chang WC et al. demonstrated that additional exercise training, which emphasizes vestibular stimulation, can improve balance ability and functional gait performance among patients with benign paroxysmal positional vertigo of the posterior semicircular canal who had already undergone the canalith repositioning manoeuvre (Chang, et al., 2008).

Vestibular rehabilitation is thought to improve long-term outcomes for BPPV. Although data are mixed, a few studies have indicated that use of vestibular rehabilitation may decrease recurrence rates for BPPV (Angeli, et al., 2003) (Helminski, et al., 2005). This protective effect against recurrence of vestibular rehabilitation may be more pronounced in the elderly (Angeli, et al., 2003). Several prospective studies have demonstrated the safety and effectiveness of vestibular rehabilitation for unilateral peripheral vestibular disorders; the results are summarized in a recent Cochrane collaboration report (Hillier, et al., 2007). Among 21 included randomized trials, there were no reports of adverse effects due to vestibular rehabilitation therapy. Current published evidence is inadequate to indicate superiority for one form of vestibular rehabilitation vs another. There is also not enough evidence to favor formal outpatient vestibular therapy performed with a clinician over independent home therapy (Kammerlind, et al., 2005).

In summary, with respect to posterior canal BPPV, vestibular rehabilitation demonstrates superior treatment outcomes compared with placebo. In short-term evaluation, vestibular rehabilitation is less effective at producing complete symptom resolution than PRMs. With longer-term follow-up, however, its effectiveness approaches that of PRMs. Insufficient data exist concerning the response of lateral canal BPPV to vestibular therapy; this area needs further research.

Cost considerations may become important if repeated visits for clinician-supervised therapy are required as opposed to initial patient instruction followed by home-based therapy. Patients with certain comorbidities may not be appropriate candidates for vestibular rehabilitation or may need specialized, individually tailored vestibular rehabilitation protocols. Examples of such comorbidities include cervical stenosis, Down syndrome, severe rheumatoid arthritis, cervical radiculopathies, Paget's disease, morbid obesity, ankylosing spondylitis, low back dysfunction, and spinal cord injuries. On the other hand, patients with preexisting otological or neurological disorders may derive more benefit from vestibular rehabilitation as a treatment for BPPV.

Verantwoording

Laatst beoordeeld : 01-02-2020

Laatst geautoriseerd : 01-02-2020

Voor de volledige verantwoording, evidence tabellen en eventuele aanverwante producten raadpleegt u de Richtlijndatabase.

Referenties

- Angeli, S.I., Hawley, R., Gomez, O. (2003). Systematic approach to benign paroxysmal positional vertigo in the elderly. *Otolaryngol Head Neck Surg*, 128, 719-25.
- Banfield, G.K., Wood, C., Knight, J. (2000). Does vestibular habituation still have a place in the treatment of benign paroxysmal positional vertigo?. *J Laryngol Otol*, 114, 501-5.
- Brandt, T., Steddin, S., Daroff, R.B. (1994). Therapy for benign paroxysmal positioning vertigo, revisited. *Neurology*, 44, 796-800.
- Brandt, T., Daroff, R.B. (1980). Physical therapy for benign paroxysmal positional vertigo. *Arch Otolaryngol*, 106, 484-5.
- Cawthorne, T. (1944). The physiologic basis for head exercises. *J Chart Soc Physiother*, 0, 106-7.
- Chang, W.C., Yang, Y.R., Hsu, L.C., Chern, C.M., Wang, R.Y. (2008). Balance improvement in patients with benign paroxysmal positional vertigo. *Clinical rehabilitation Apr*, 22 (4), 338-47.
- Cohen, H.S., Kimball, K.T. (2005). Effectiveness of treatments for benign paroxysmal positional vertigo of the posterior canal. *Otol Neurotol*, 26, 1034-40.
- Furman, J.M., Cass, S.P. (1999). Benign paroxysmal positional vertigo. *N Engl J Med*, 341, 1590-6.
- Helminski, J.O., Janssen, I., Kotaspouikis, D., et al. (2005). Strategies to prevent recurrence of benign paroxysmal positional vertigo. *Arch Otolaryngol Head Neck Surg*, 131, 344-8.
- Herdman, S.J., Blatt, P.J., Schubert, M.C. (2000). Vestibular rehabilitation of patients with vestibular hypofunction or with benign paroxysmal positional vertigo. *Curr Opin Neurol*, 13, 39-43.
- Hillier, S.L., Hollohan, V. (2007). Vestibular rehabilitation for unilateral peripheral vestibular dysfunction. *Cochrane Database Syst*, 0, CD005397-.
- Kammerlind, A.S., Ledin, T.E., Odkvist, L.M., et al. (2005). Effects of home training and additional physical therapy on recovery after acute unilateral vestibular loss—a randomized study. *Clin Rehabil*, 19, 54-62.
- Norré, M.E., Forrez, G., Beckers, A. (1987). Vestibular habituation training and posturography in benign paroxysmal positioning vertigo. *ORL J Otorhinolaryngol Relat Spec.*, 49 (1), 22-5.
- Norré, M.E., Beckers, A. (1987). Exercise treatment for paroxysmal positional vertigo: comparison of two types of exercises. *Arch Otorhinolaryngol*, 244 (5), 291-4.
- Steenerson, R.L., Cronin, G.W. (1996). Comparison of the canalith repositioning procedure and vestibular habituation training in forty patients with benign paroxysmal positional vertigo. *Otolaryngol Head Neck Surg*, 114, 61-4.
- Soto Varela, A., Bartual Magro, J., Santos Perez, S., et al. (2001). Benign paroxysmal vertigo: a comparative prospective study of the efficacy of Brandt and Daroff exercises, Semont and Epley maneuver. *Rev Laryngol Otol Rhinol (Bord)*, 122, 179-83.
- Telian, S.A., Shepard, N.T. (1996). Update on vestibular rehabilitation therapy. *Otolaryngol Clin North Am*, 29, 359-71.
- Toledo, H., Cortes, M.L., Pane, C., et al. (2000). Semont maneuver and vestibular rehabilitation exercises in the treatment of benign paroxysmal postural vertigo. A comparative study. *Neurologia*, 15, 152-7.
- Troost, B.T., Patton, J.M. (1992). Exercise therapy for positional vertigo. *Neurology*, 42, 1441-4.
- Whitney, S.L., Rossi, M.M. (2000). Efficacy of vestibular rehabilitation. *Otolaryngol Clin North Am*, 33, 659-72.

Medicatie als behandeling van BPPD

Uitgangsvraag

Zijn medicijnen geschikt als therapie om patiënten met BPPD te behandelen?

Aanbeveling

Er is geen indicatie om patiënten met BPPD medicatie voor te schrijven.

Overwegingen

- Voordeel: patiënt krijgt geen onterechte medicatie
- Nadeel: geen
- Kosten: voordelig om geen medicatie voor te schrijven
- Afweging: voordelen wegen op tegen de nadelen. Kortdurend gebruik van vestibulosuppressieve medicatie en/of anti-emetica kan soms wel zinvol zijn om een Dix-Hallpikemanoeuvre of repositiemanoeuvre te kunnen uitvoeren teneinde misselijkheid en/of braken te voorkomen.
- Waarde oordeel: Schade door ineffectieve behandeling wordt voorkomen.
- Rol van de voorkeur van de patiënt: is minimaal.
- Exclusiecriteria: Patiënten die profylaxe voor Dix-Hallpikemanoeuvre en/of repositie manoeuvre nodig hebben.

Onderbouwing

Conclusies

Niveau 3	Medicatie is niet effectief als behandeling van BPPD.
-----------------	---

Samenvatting literatuur

The symptoms of vertigo due to many different underlying etiologies are commonly treated with medications. Clinicians may prescribe pharmacological management to either 1) reduce the spinning sensations of vertigo specifically and/or 2) to reduce the accompanying motion sickness symptoms. These motion sickness symptoms include a constellation of autonomic or vegetative symptoms such as nausea, vomiting, and diarrhea, which can accompany the vertigo. Such pharmacological therapies for vertigo may be broadly termed *vestibular suppressant medications* (Hain, et al., 2003) (Hain, et al., 2005). Several categories of vestibular suppressant medications are in common use. Of these, the most commonly used are benzodiazepines and antihistamines. Benzodiazepines, such as diazepam and clonazepam, have anxiolytic, sedative, muscle relaxant, and anticonvulsant properties derived from potentiating the inhibitory effect of the gamma-amino butyric acid system. In prolonged dizziness, these medications can reduce the subjective sensation of spinning, but they also interfere with central compensation in peripheral vestibular conditions. Antihistamines, on the other hand, appear to have a suppressive effect on the central emetic center to relieve the nausea and vomiting associated with motion sickness. Common examples of antihistamines used to treat symptoms of vertigo and/or associated motion sickness include meclizine and diphenhydramine. Other medications that are often used for

motion sickness include promethazine, which is a phenothiazine with antihistamine properties, and ondansetron, which is a serotonin-5-hydroxytryptamine-3 antagonist. Finally, anticholinergic medications such as scopolamine block acetylcholine, which is a widespread CNS transmitter, and help with motion sickness by reducing neural mismatching (Hain et al., 2003) (Hain, et al., 2005).

There is no evidence in the literature to suggest that any of these vestibular suppressant medications are effective as a definitive, primary treatment for BPPD, or as a substitute for repositioning maneuvers (Frohman, et al., 2003) (Hain, et al., 2003) (Carlow, et al., 1986) (Cesarani, et al., 2004) (Fujino, et al., 1994). Exercise was found to be a better treatment choice than medication (betahistin) and may be preferable for patients with persistent or chronic vertigo (Kulcu, et al., 2008). Some studies show a resolution of BPPD over time with medications, but these studies follow patients for the period of time in which spontaneous resolution would occur (Woodworth, et al., 2004) (Salvinelli, et al., 2004) (Itaya, et al., 1997) (McClure, et al., 1980). In one double-blind controlled trial by McClure and Willet (McClure, et al., 1980) comparing diazepam, lorazepam, and placebo, all groups showed a gradual decline in symptoms with no additional relief in the drug treatment arms. In a small study, Itaya et al (Itaya, et al., 1997) compared PRMs to a medication-alone treatment arm and found that PRMs had substantially higher treatment responses (78.6%-93.3% improvement) compared with medication alone (30.8% improvement) at 2 weeks follow-up. These data reinforced previous data from Fujino et al (Fujino, et al., 1994) that also indicated superiority of vestibular training for BPPD over medication use alone. A lack of benefit from vestibular suppressants and their inferiority to PRMs indicate that clinicians should not substitute pharmacological treatment of symptoms associated with BPPD in lieu of other more effective treatment modalities.

Conversely, vestibular suppressant medications have the potential for significant harm. All of these medications may produce drowsiness, cognitive deficits, and interference with driving vehicles or operating machinery (Ancelin, et al., 2006) (Hebert, et al., 2007) (Barbone, et al., 1998) (Engeland, et al., 2007) (Jauregui, et al., 2006). Medications used for vestibular suppression, especially psychotropic medications such as benzodiazepines, are a significant independent risk factor for falls (Hartikainen, et al., 2007). The risk of falls increases in patients taking multiple medications and with the use of medications such as antidepressants (Lawson, et al., 2005) (Hien, et al., 2005). The potential for polypharmacy when adding vestibular suppressants further exposes the elderly to additional risk (Landi, et al., 2007). Educational programs to modify practitioner's use of such medications can result in a reduction of falls (Pit, et al., 2007).

There are other potential harmful side effects of vestibular suppressants. Benzodiazepines and antihistamines interfere with central compensation for a vestibular injury (Hanley, et al., 1998) [Baloh, 1998] (Baloh, et al., 1998). The use of vestibular suppressants may obscure the findings on the Dix-Hallpike maneuvers. In addition, there is evidence of additional potential harm from the antihistamine class of medications on cognitive functioning (Ancelin, et al., 2006), and on gastrointestinal motility, urinary retention, vision, and dry mouth in the elderly (Rudolph, et al., 2008).

Another type of medication, betahistine dihydrochloride is an extensively applied and studied drug in the treatment of vertigo as well, especially in case of Meniere Disease. Betahistine appears to be a weak H1 agonist (release of Ca^{2+}), a weak H2 agonist (synthesis cyclic AMP) and a strong H3 antagonist (release of histamine) (Timmerman, et al., 1989). The action upon the H3 auto-receptor might explain why a relatively low

concentration of betahistine could effectively modify neurotransmission in the brain (Tighilet, et al., 1995). Betahistine also results in a dose-dependent inhibition of polysynaptic neurons in the lateral vestibular nuclei (KawabataA). Recently it was shown that betahistine fastens the central compensation process after labyrinthectomy in cats (Tighilet, et al., 1995) and after neurectomy in men (Redon et al., 2010), In animals intralabyrinthine blood flow and oxygenation of sensory tissue is increased by intake of betahistine (Meyer et al., 1994, Laurikainen et al., 1993). In one study, the treatment of patients with BPPV was found not to be effective in BPPD (Kulcu, et al., 2008).

In summary, vestibular medications are not recommended for treatment of BPPD, other than for the short-term management of vegetative symptoms such as nausea or vomiting in a severely symptomatic patient. Examples of potential short-term uses include patients who are severely symptomatic yet refuse therapy or patients who become severely symptomatic after a PRM. Antiemetics may also be considered for prophylaxis for patients who have previously manifested severe nausea and/or vomiting with the Dix-Hallpike maneuvers and in whom a PRM is planned. If prescribed for these very specific indications, clinicians should also provide counseling that the rates of cognitive dysfunction, falls, drug interactions, and machinery and driving accidents increase with use of vestibular suppressants.

Verantwoording

Laatst beoordeeld : 01-02-2020

Laatst geautoriseerd : 01-02-2020

Voor de volledige verantwoording, evidence tabellen en eventuele aanverwante producten raadpleegt u de Richtlijndatabase.

Referenties

- Ancelin, M.L., Artero, S., Portet, F., et al. (2006). Non-degenerative mild cognitive impairment in elderly people and use of anticholinergic drugs: longitudinal cohort study. *BMJ*, 332, 455-9.
- Baloh, R.W. (1998). Vertigo. *Lancet*, 352, 1841-6.
- Baloh, R.W. (1998). Dizziness: neurological emergencies. *Neurol Clin*, 16, 305-21.
- Barbone, F., McMahon, A.D., Davey, P.G., et al. (1998). Association of roadtraffic accidents with benzodiazepine use. *Lancet*, 352, 1331-6.
- Carlow, T.J. (1986). Medical treatment of nystagmus and ocular motor disorders. *Int Ophthalmol Clin*, 26, 251-64.
- Cesarani, A., Alpini, D., Monti, B., et al. (2004). The treatment of acute vertigo. *Neurol Sci*, 25, S26-S30.
- Engeland, A., Skurtveit, S., Morland, J. (2007). Risk of road traffic accidents associated with the prescription of drugs: a registry-based cohort study. *Ann Epidemiol*, 17, 597-602.
- Frohman, E.M., Kramer, P.D., Dewey, R.B., et al. (2003). Benign paroxysmal positioning vertigo in multiple sclerosis: diagnosis, pathophysiology and therapeutic techniques. *Mult Scler*, 9, 250-5.
- Fujino, A., Tokumasu, K., Yosio, S., et al. (1994). Vestibular training for benign paroxysmal positional vertigo. Its efficacy in comparison with antivertigo drugs. *Arch Otolaryngol Head Neck Surg*, 120, 497-504.
- Hain, T.C., Yacovino, D. (2005). Pharmacologic treatment of persons with dizziness. *Neurol Clin*, 23, 831-53.
- Hain, T.C., Uddin, M. (2003). Pharmacological treatment of vertigo. *CNS Drugs*, 17, 85-100.
- Hanley, K., O'Dowd, T., Considine, N. (2001). A systematic review of vertigo in primary care. *Br J Gen Pract*, 51, 666-71.
- Hartikainen, S., Lonroos, E., Louhivuori, K. (2007). Medication as a risk factor for falls: critical systematic review. *J Gerontol A Biol Sci Med Sci*, 62, 1172-81.
- Hebert, C., Delaney, J.A., Hemmelgarn, B., et al. (2007). Benzodiazepines and elderly drivers: a comparison of pharmacoepidemiological study designs. *Pharmacoepidemiol Drug Saf*, 16, 845-9.

- Hien le, T.T., Cumming, R.G., Cameron, I.D., et al. (2005). Atypical antipsychotic medications and risk of falls in residents of aged care facilities. *J Am Geriatr Soc*, 53, 1290-5.
- Itaya, T., Yamamoto, E., Kitano, H., et al. (1997). Comparison of effectiveness of maneuvers and medication in the treatment of benign paroxysmal positional vertigo. *ORL J Otorhinolaryngol Relat Spec*, 59, 155-8.
- Jauregui, I., Mullol, J., Bartra, J., et al. (2006). H1 antihistamines: psychomotor performance and driving. *J Investig Allergol Clin Immunol*, 16, 37-44.
- Kulcu, D.G., Yanik, B., Boynukalin, S., Kurtais, Y. (2008). Efficacy of a home-based exercise program on benign paroxysmal positional vertigo compared with betahistine. *J Otolaryngol Head Neck Surg*. Jun, 37 (3), 373-9.
- Landi, F., Russo, A., Liperoti, R., et al. (2007). Anticholinergic drugs and physical function among frail elderly population. *Clin Pharmacol Ther*, 81, 235-41.
- Laurikainen, E.A., Miller, J.M., Quirk, W.S., Kallinen, J., Ren, T., Nuttall, A.L., Grenman, R., Virolainen, E. (1993). Betahistine induced vascular effects in the rat cochlea. *Am J Otol*. Jan, 14 (1), 24-30.
- Lawson, J., Johnson, I., Bamiou, D.E., et al. (2005). Benign paroxysmal positional vertigo: clinical characteristics of dizzy patients referred to a Falls and Syncope Unit. *QJM*, 98, 357-64.
- McClure, J.A., Willett, J.M. (1980). Lorazepam and diazepam in the treatment of benign paroxysmal vertigo. *J Otolaryngol*, 9, 472-7.
- Meyer, P., Schmidt, R., Grutzmacher, W., Gehrig, W. (1994). Inner ear blood flow with betahistine an animal experiment study. *Laryngorhinootologie*, 73 (3), 153-6.
- Pit, S.W., Byles, J.E., Henry, D.A., et al. (2007). A Quality Use of Medicines program for general practitioners and older people: a cluster randomized controlled trial. *Med J Aust*, 187, 23-30.
- Redon, C., Lopez, C., Bernard-Demanze, L., Dumitrescu, M., Magnan, J., Lacour, M., Borel, L. (2010). Betahistine Treatment Improves the Recovery of Static Symptoms in Patients With Unilateral Vestibular Loss. *J Clin Pharmacol*, 0, X.
- Rudolph, J.L., Salow, M.J., Angelini, M.C., et al. (2008). The anticholinergic risk scale and anticholinergic adverse effects in older persons. *Arch Intern Med*, 168, 508-13.
- Salvinelli, F., Trivelli, M., Casale, M., et al. (2004). Treatment of benign positional vertigo in the elderly: a randomized trial. *Laryngoscope*, 114, 827-31.
- Tighilet, B., Leonard, J., Lacour, M. (1995). Betahistine dihydrochloride treatment facilitates vestibular compensation in the cat. *J Vestib Res*, 5 (1), 53-66.
- Timmerman, H. (1989). The Histamine H3-receptor, its function and ligands. In van der Goot, H., Pallos, L., Timmerman, H., eds. *Trends in medicinal chemistry '88*. Amsterdam, 0, 351-63.
- Woodworth, B.A., Gillespie, M.B., Lambert, P.R. (2004). The canalith repositioning procedure for benign positional vertigo: a meta-analysis. *Laryngoscope*, 114, 1143-6.

Chirurgische interventie als behandeling van BPPD

Uitgangsvraag

Is chirurgische interventie geschikt als behandeling?

Aanbeveling

De werkgroep beveelt aan om bij ernstig invaliderende, therapieresistente, BPPD canal plugging te overwegen. De kans op ernstige bijwerkingen zoals doofheid en blijvende evenwichtsstoornissen dient goed met de patiënt te worden besproken.

Neurectomie van de n. singularis is een techniek die gezien de complexiteit voorbehouden is aan gespecialiseerde KNO-artsen.

Overwegingen

- Voordeel: bewezen effectiviteit
- Nadelen: kans op perceptieslechthoerendheid, kans op (blijvende) evenwichtsstoornissen. Neurectomie is een technisch veeleisende operatie waarbij de ervaring van de chirurg een grote rol speelt.
- Kosten: fors
- Afweging van voordeel tegen nadeel: bij therapieresistente BPPD moeten de risico's van een operatieve ingreep worden afgewogen tegen de ernst van de klachten/invalidering door de klachten
- Waarde oordelen:
- Rol van de voorkeur van de patiënt: groot
- Exclusie: geen

Onderbouwing

Achtergrond

Benign paroxysmal positional vertigo (BPPV) is one of the most common disorders in patients suffering from vertigo. The pathogenesis is currently thought to be canalolithiasis or cupulolithiasis of the posterior semicircular canal (Epley, et al., 1990). Canalith repositioning procedures by which densities are removed from the responsible canal are widely accepted in the treatment of patients with BPPV (Epley, et al., 1992; Seo et al., 2007). Positional vertigo is immediately resolved by the treatment; however, a few patients continue to complain of vertigo after treatment (Seo et al., 2007).

According to the canalolithiasis theory, the symptoms should improve when the endolymphatic current of the responsible canal is blocked. Parnes and McClure were the first to report that positional vertigo completely disappeared after posterior semicircular canal occlusion surgery in two cases of intractable BPPV in 1990 (Parnes, McClure, et al., 1990 4). Although later studies also showed that plugging surgery has a high success rate for resolving positional vertigo and positional nystagmus, it is not widely performed. One reason is a risk of inner ear injury that causes severe sensorineural hearing loss (SNHL) and prolonged disequilibrium.

Conclusies

Niveau 3	<p>Het lijkt waarschijnlijk dat canal plugging een effectieve en veilige behandelmethodede voor BPPD is als herhaalde repositiemanoeuvres niet werken.</p> <p><i>bronnen niveau D: Fife et.al., 2008, Agrawal and Parnes, et al., 2005, Seo et.al., 2009</i></p>
----------	--

Niveau 3/4	<p>Er zijn aanwijzingen dat neurectomie van de n. singularis effectief is bij de behandeling van BPPD, maar deze ingreep gaat gepaard met een grote kans op bijwerkingen zoals perceptief gehoorsverlies.</p> <p><i>bronnen niveau D: (Gacek et al.,), (Schessel, et al., 1998), (Leveque, et al., 2007)</i></p>
------------	--

Samenvatting literatuur

Canal plugging

One evidence-based review was found that addressed the question whether surgical occlusion of the posterior canal or singular neurectomy is effective for BPPV (Fife et.al., 2008). The most common used procedure in this studies is fenestration and occlusion of the posterior semicircular canal. Five studies of level D, including in total 86 patients undergoing canal occlusion, reported 'complete relief' of BPPV symptoms in 85 as ascertained by the treating surgeon. Reported complications included a 'mild' conductive hearing loss for 4 weeks or less, 'mild' and transient' unsteadiness in most patients, and a high frequency sensorineural hearing loss in 6 patients. Two other level D studies were found that used canal plugging to treat BPPV patients. One study described results of 48 patients. A limitation of this study was that inclusion criteria for surgery were not clearly described. It was not clear whether these patients suffered from intractable BPPV. Out of 44 patients with normal preoperative hearing, one had a delayed (3-months) sudden and permanent profound loss, while another had a mild (20dB) loss. Six patients had protracted courses of imbalance and motion sensitivity (Agrawal and Parnes, et al., 2005). The other study specifically looked into the effect of posterior semicircular canal plugging surgery on hearing and balance functions. The patients included in this study were five consecutive patients with intractable BPPV, 1% of all BPPV patients treated in this period, who underwent plugging surgery. After surgery positional vertigo was resolved in all patients. This study showed that results of audiometry, caloric testing and vestibular evoked myogenic potential (VEMP) testing were hardly influenced by plugging surgery (Seo et.al., 2009).

Singular neurectomy

Singular neurectomy is another surgical treatment for intractable BPPV. It relieves the positional vertigo by deafferentiating the ampulla of the posterior semicircular canal. In this procedure, the singular canal in the round window niche is identified and the nerve transected. In one study 96.8% were reported to have 'complete relief' of vertigo after singular neurectomy (Gacek, et al., 1974).

Further reports by Gacek demonstrated high efficacy with complete vertigo resolution in 91.7% of patients (Gacek, et al., 1978). A non-systematic review (Leveque et al., 2007) included five other studies that performed singular neurectomy. Complete relief was seen in 75% to 96% of cases according to the test performed by the surgeon. Partial relief occurred in 1.5% to 17% of patients, which is defined by reduction but not absence of vertigo and nystagmus triggered by a provocative maneuver. There were some patients whose symptoms did

not improve after surgery, mostly because of incorrect diagnoses, failure to locate the singular nerve, and incomplete transection of the posterior ampullary nerve due to the presence of an accessory branch. Another study, not included in this review, evaluated eight patients with BPPV, one with typical symptoms, but without nystagmus at the Hallpike's manoeuvre (Pournaras et al., 2008). After singular neurectomy all patients were free of vertigo, but sensorineural hearing loss occurred in two patients.

Sensorineural hearing loss depended on the surgeon's experience. Gacek et al. reported an initial 7.3% risk of sensorineural hearing loss with the procedure. This risk was decreased in an update of his personal series in 1995 to 3%, for 252 neurectomies (Gacek 1995; Agrawal et al., 2005). Other reported a risk of hearing loss that ranged from 9% (Silverstein, et al., 1990), 19% (Meyerhoff, et al., 1985) to 41% of cases for Epley (12 neurectomies) (Epley, et al., 1980). It ranges from a 30 dB hearing loss to total deafness. Most of those sensorineural hearing losses can be explained by the surgical act (injury to the scala tympani, to the perilymphatic space or, worse, to the endolymphatic space). Sensorineural hearing losses that remain unexplained may be due to secondary labyrinthitis.

Patients experienced dizziness for a few days after the intervention; for that period, a vertical downbeat nystagmus can be seen that is known to be of cerebellar origin. Silverstein and White explored the vestibular function on operated patients. They found that 41% of them had vestibular dysfunction with no clinical consequences whereas there were only 14% before operation (Leveque et al., 2007).

Comparison with canal occlusion:

Although singular neurectomy seems safe in experienced hands, it is technically very demanding with a high risk of sensorineural hearing loss (Schessel, et al., 1998). This may explain why it has largely been replaced by the simpler semicircular canal occlusion.

Verantwoording

Laatst beoordeeld : 01-02-2020

Laatst geautoriseerd : 01-02-2020

Voor de volledige verantwoording, evidence tabellen en eventuele aanverwante producten raadpleegt u de Richtlijndatabase.

Referenties

- Agrawal, S.K., Parnes, L.S. (2005). Surgical treatment of benign paroxysmal positional vertigo. *Audiological Medicine*, 3 (1), 63-68.
- Epley, J.M. (1990). New dimensions of benign paroxysmal positional vertigo. *Otolaryngol Head Neck Surg*, 88, 599-605.
- Epley, J.M. (1992). The canalith repositioning procedure: for treatment of benign paroxysmal positional vertigo. *Otolaryngol Head Neck Surg*, 107, 399-404.
- Epley, J.M. (1980). Singular neurectomy: hypotympanotomy approach. *Otolaryngol Head Neck Surg*, 88, 304-9.
- Fife, T.D., Iverson, D.J., Lempert, T., et al. (2008). Practice parameter: therapies for benign paroxysmal positional vertigo (an evidence-based review): Report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology*, 70, 2067-74.
- Gacek, R.R. (1978). Singular neurectomy update. *Ann Otol Rhinol Laryngol*, 87, 300-305.
- Gacek, R.R. (1995). Technique and results of singular neurectomy for the management of benign paroxysmal positional vertigo. *Acta Otolaryngol*, 115, 154-7.

- Leveque, M., Labrousse, M., Seidermann, L., Chays, A. (2007). Surgical therapy in intractable benign paroxysmal positional vertigo. *Otolaryngology - Head & Neck Surgery*, 136 (5), 693-8.
- Meyerhoff, W.L. (1985). Surgical section of the posterior ampullary nerve. *Laryngoscope*, 95, 933-935.
- Parnes, L.S., McClure, J.A. (1990). Posterior semicircular canal occlusion for intractable benign paroxysmal positional vertigo. *Ann Otol Rhinol Laryngol*, 99, 330-4.
- Pournaras, I., Kos, I., Guyot, J.P. (2008). Benign paroxysmal positional vertigo: a series of eight singular neurectomies. *Acta Oto-Laryngologica*, 128 (1), 5-8.
- Schessel, D.A., Minor, L.B., Nedzelski, J.M. (1998). Meniere's disease and other peripheral vestibular disorders. In: Cummings C, editor . *Otolaryngology – Head & Neck Surgery*, Mosby, 0, X-.
- Seo, T., Hashimoto, M., Saka, N., Sakagami, M. (2009). Hearing and vestibular functions after plugging surgery for the posterior semicircular canal. *Acta Oto-Laryngologica*, 129, 1148-1152.
- Seo, T., Miyamoto, A., Saka, N., Shimano, K., Sakagami, M. (2007). Immediate efficacy of the canalith repositioning procedure for the treatment of benign paroxysmal positional vertigo. *Otol Neurotol*, 28, 917-9.
- Silverstein, H., White, D.W. (1990). Wide surgical exposure for singular neurectomy in the treatment of benign positional vertigo. *Laryngoscope*, 100, 701-706.

Expectatief beleid bij BPPD

Uitgangsvraag

Kan bij patiënten met BPPD volstaan worden met een expectatief beleid?

Aanbeveling

Behandelen van BPPD verdient de voorkeur boven het afwachten van het natuurlijk beloop.

Overwegingen

- Voordeel: De symptomen verdwijnen bij 35-50 procent van de patiënten na ongeveer 1 maand zonder interventie.
- Nadeel: Langer last van symptomen dan bij behandeling waardoor de patiënt een groter valrisico heeft of niet in staat is tot werken.
- Kosten: indirecte kosten vanwege een vertraagde genezing in vergelijking met andere behandelingen.
- Afweging: Een afwachtend beleid als behandeling voor posterieur kanaal BPPD heeft als voordeel dat de patiënt geen manoeuvres of vestibulaire revalidatie hoeft te ondergaan. Niet behandelen heeft echter als nadeel dat de klachten langer persisteren met mogelijke gevolgen voor kwaliteit van leven, vallen en arbeidsverzuim..
- Waarde oordeel: de werkgroep heeft een voorkeur voor een behandelingsinterventie in plaats van het natuurlijk beloop af te wachten, vooral omdat de klachten dan sneller verdwijnen.
- Rol van de voorkeur van de patiënt: substantieel gezien de gezamenlijke beslissing over de behandeling.
- Exclusiecriteria: geen

Onderbouwing

Conclusies

Niveau 2	De symptomen verdwijnen bij 35-50 procent van de patiënten na ongeveer 1 maand zonder interventie.
-----------------	--

Samenvatting literatuur

Observation may be defined as a “watchful waiting” or the withholding of specific therapeutic interventions for a given disease. Observation is often considered when the disease course is self-limited and/or felt to be benign with limited sequelae occurring from the withholding of therapy. In BPPV, observation implies that therapeutic interventions such as vestibular rehabilitation and/or PRMs will be withheld, anticipating a natural and spontaneous improvement of the symptoms of BPPV. Under a course of observation, patients may still be instructed to avoid provocative positions and activities where the risk of injury (ie, falls) may be increased until symptoms resolve spontaneously or until they are reassessed for symptom resolution.

To consider observation as an option in the management of BPPV, the clinician must determine the natural history of the BPPV. It has been presumed that the natural history of BPPV is one of eventual resolution in most

patients. It should be noted, however, that an often quoted study by Blakley, (Blakley, et al., 1994) which reported high rates of spontaneous resolution of BPPV, relied on subjective symptom reporting, rather than objective testing with a Dix-Hallpike maneuver, as the outcome measure for resolution. It is believed that a significant fraction of patients reporting subjective improvement actually have reduction in symptoms secondary to avoiding provocative (vertigo-producing) positions rather than actual cure (Woodworth, et al., 2004). More recent RCTs have utilized objective testing with the Dix-Hallpike maneuver as an additional outcome measure to assess for objective resolution of BPPV. Notably, to observe proper blinding, most RCTs also use a sham positional maneuver in the control group, which theoretically may affect the natural history of BPPV.

In several studies, the spontaneous rate of symptomatic resolution of BPPV ranges from 15 to 86 percent. The reported rate of spontaneous improvement based on objective positional testing (ie, conversion to a negative Dix- Hallpike maneuver) ranges from 35 percent to 50 percent (Woodworth, et al., 2004). As demonstrated in Table 4.1, the natural history of posterior canal BPPV varies widely across studies at a 1-month and a 3-month follow-up interval. Further variability in the spontaneous resolution rate arises from differences in duration of symptoms prior to actual diagnoses of BPPV as well as differences in duration of follow-up (Hilton, et al., 2004) (Froehling, et al., 2000) (Lynn, et al., 1995) (Sekine, et al., 2006). Longitudinal follow-up studies of untreated BPPV patients are lacking, but one study of completely untreated patients determined a mean time interval from onset of symptoms to spontaneous resolution of BPPV of 39 _ 47 days (Imai, et al., 2005). As would be expected, spontaneous symptom resolution rates increase with increasing duration of follow-up among observed patients.

Tabel 4.1: Symptoom reductie rates voor observatie alleen van BPPD

Referentie	Genezen n/m	% genezen	Placebo behandeling of pure observatie	Tijd tot assessment
Von Brevern, 2007	22/26	84.6%	Placebo	4 weken
Sakine 2006	48/60	80.0%	Observatie	1 maand
Imai 2005	45/70	64.0	Observatie	1 maand
Simhadri 2003	3/15	20	Observatie	4 weken
Yimtae 2003	7/20	35.0	Observatie	1 maand
Sherman 2001	11/22	50.0	Placebo	3 maanden
Asawavichianginda 2000	18/22	81.8	Observatie	3 maanden
Steenerson 1996	17/40	42.5	Observatie	3 maanden
Lynn 1995	3/15	20.0	Placebo	1 maand
Blakley 1994	19/22	86.4	Observatie	1 maand

Eindpunt: herstel van vertigo symptomen op het tijdstip van assessment.

Although observation of posterior canal BPPV is an option for management, clinicians should also be aware that other treatments such as the PRM have been shown to offer patients faster resolution of BPPV symptoms. A meta-analysis of nine separate trials examining the efficacy of the PRM for BPPV treatment demonstrated

consistent improvement in the treatment group, with up to 4.1 times greater rates of symptom resolution (95% confidence interval, 3.1- 5.2) in the PRM groups vs the control groups at initial assessments within 1 month. Studies with follow-up at beyond 1 month still demonstrated an improvement rate of nearly three times that of controls (Woodworth, et al., 2004). Other longer-term follow-up data also suggest that patients treated with a PRM had lower rates of relapse of BPPV at 6 months and 1 year posttreatment (Simhadri, et al., 2003).

Observation as an option for the management of posterior canal BPPV offers the potential benefits of avoiding repositioning maneuvers or vestibular rehabilitation, which in turn may provoke symptoms and discomfort. There may also be a cost savings from decreased rates of referral for vestibular rehabilitation or PRMs. From a potential harms perspective, patients who elect for the observation option should be informed about a typically longer duration of symptoms compared with a treatment maneuver and potentially higher recurrence rates. Appropriate precautions for the risks associated with BPPV symptoms should be taken during the watchful waiting period.

The natural history of lateral canal BPPV is less well defined than that of posterior canal BPPV. Several authors have commented that lateral canal BPPV may be prone to more rapid spontaneous resolution than posterior canal BPPV (Moon, et al., 2006) (Sekine, et al., 2006). In one study, the mean time between the onset of vertigo in lateral canal BPPV to spontaneous resolution was 16 _ 19 days (Imai, et al., 2005). Although repositioning maneuvers have shown success in lateral canal BPPV, overall high quality comparative data regarding treatment vs observation such as RCTs are limited in this subtype of BPPV (Casani, et al., 2002) (Sekine, et al., 2006) (Fife, et al., 2006). Thus, observation of lateral canal BPPV remains an option for management. Future RCTs need to be dedicated to the interventional management of lateral canal BPPV.

Verantwoording

Laatst beoordeeld : 01-02-2020

Laatst geautoriseerd : 01-02-2020

Voor de volledige verantwoording, evidence tabellen en eventuele aanverwante producten raadpleegt u de Richtlijndatabase.

Referenties

- Blakley, B.W. (1994). A randomized, controlled assessment of the canalith repositioning maneuver. *Otolaryngol Head Neck Surg*, 110, 391-6.
- Casani, A.P., Vannucci, G., Fattori, B., et al. (2002). The treatment of horizontal canal positional vertigo: our experience in 66 cases. *Laryngoscope*, 112, 172-8.
- Fife, T.D. (1998). Recognition and management of horizontal canal benign positional vertigo. *Am J Otol*, 19, 345-51.
- Eroehling, D.A., Bowen, J.M., Mohr, D.N., et al. (2000). The canalith repositioning procedure for the treatment of benign paroxysmal positional vertigo: a randomized controlled trial. *Mayo Clin Proc*, 75, 695-700.
- Hilton, M., Pinder, D. (2004). The Epley (canalith repositioning) manoeuvre for benign paroxysmal positional vertigo. *Cochrane Database Syst Rev*, 0, CD003162-.
- Imai, T., Ito, M., Takeda, N., et al. (2005). Natural course of the remission of vertigo in patients with benign paroxysmal positional vertigo. *Neurology*, 64, 920-1.
- Lynn, S., Pool, A., Rose, D., et al. (1995). Randomized trial of the canalith repositioning procedure. *Otolaryngol Head Neck Surg*, 113, 712-20.
- Moon, S.Y., Kim, J.S., Kim, B.K., et al. (2006). Clinical characteristics of benign paroxysmal positional vertigo in Korea: a

multicenter study. *J Korean Med Sci*, 21, 539-43.

Sekine, K., Imai, T., Sato, G., et al. (2006). Natural history of benign paroxysmal positional vertigo and efficacy of Epley and Lempert maneuvers. *Otolaryngol Head Neck Surg*, 135, 529-33.

Simhadri, S., Panda, N., Raghunathan, M. (2003). Efficacy of particle repositioning maneuver in BPPV: a prospective study. *Am J Otolaryngol*, 24, 355-60.

Woodworth, B.A., Gillespie, M.B., Lambert, P.R. (2004). The canalith repositioning procedure for benign positional vertigo: a meta-analysis. *Laryngoscope*, 114, 1143-6.

Omgevingsfactoren die belangrijk zijn voor de behandeling van BPPD

Uitgangsvraag

Met welke factoren moet rekening worden gehouden bij de behandeling van BPPD?

Aanbeveling

Artsen dienen factoren die een verhoogd risico op vallen geven uit te vragen; dit beïnvloedt de behandelkeus van BPPD (voorkeur voor niet-conservatieve behandeling).

Overwegingen

- Voordeel: Het behandelplan voor BPPD wordt patiënt-specifiek en patiënten met een verhoogd risico op vallen en valgerelateerde morbiditeit worden geïdentificeerd.
- Nadeel: geen
- Kosten: geen
- Rol van de voorkeur van de patiënt: minimaal

Onderbouwing

Conclusies

Niveau 3	BPPD gaat veelal gepaard met comorbiditeit zoals diabetes, osteoporose en vasculaire problematiek. BPPD geeft een verhoogde kans op vallen en dit neemt toe bij comorbiditeit.
-----------------	--

Samenvatting literatuur

Although BPPV arises from dysfunction of the vestibular end organ, patients with BPPV often concurrently suffer from comorbidities, limitations, and risks that may affect the diagnosis and treatment outcome of BPPV. Assessment of the patient with BPPV for factors that modify management is essential for improved treatment outcomes and ensuring patient safety with an underlying diagnosis of BPPV. The majority of factors that may modify management of BPPV can be identified if the clinician questions patients for these factors and elicits a detailed history (Rubenstein, et al., 2001).

Given that BPPV occurs most commonly in the second half of the lifespan and its prevalence increases with age, patients suffering from BPPV often have medical comorbidities that may alter the management of BPPV (Lawson, et al., 2005). In cross-sectional surveys, patients with BPPV demonstrate higher rates of diabetes, history of head trauma, and anxiety (Cohen, et al., 2004). Other studies have also found higher relative rates of migraine (34% in BPPV patients vs 10% in non-dizziness control group), history of stroke (10% in BPPV patients vs 1% in controls), diabetes (14% vs 5%), and hypertension (52% vs 22%) (von Brevern, et al., 2007). Clinicians should assess patients with BPPV for these comorbidities because their presence may modify management and influence treatment outcomes in BPPV.

One of the major concerns with BPPV and vertiginous syndromes in general is the risk for falls and resultant injury (Gazzola, et al., 2006). In multiple studies concerning etiology of falls, dizziness and vertigo were deemed

the primary etiology for 13 percent of falls, compared with existing balance and gait problems (17%) and person-environment interactions (31%) (Rubenstein, et al., 2006). In a study by Oghalai, 15.9 percent of patients referred to a geriatric clinic for general geriatric evaluation had undiagnosed BPPV, and three-fourths of those with BPPV had fallen within the 3 months prior to referral. Thus, evaluation of patients with a diagnosis of BPPV should also include an assessment of risk for falls (Lawson, et al., 2005). In particular, elderly patients will be more statistically at risk for falls with BPPV. Clinicians may use various fall assessment tools to determine the patient's fall risk and appropriate precautionary recommendations (Rubenstein, et al., 2001).

As noted above, comorbid conditions that occur commonly with BPPV such as a history of stroke or diabetes should also be identified during evaluation of patients with BPPV. Patients with a history of stroke or a history of diabetes, particularly with peripheral neuropathy, may already have preexisting gait, balance, or proprioceptive deficit (Casellini, et al., 2007) (Richardson, et al., 2002) (Tilling, et al., 2006). The additional symptoms of BPPV may increase their risk for fall and injury. Patients with visual disturbances often lack the ability to correct for or compensate for a balance deficit with visual cues, and may also be at increased risk for falls. Associations between osteopenia and osteoporosis and BPPV have been reported (Vibert, et al., 2003). Patients with both osteoporosis and BPPV may be at greater risk for fractures resulting from falls related to BPPV; therefore, patients with combined osteoporosis and subsequent BPPV should be identified and monitored closely for fall and fracture risk. Examined from a different vantage point, patients with a history of recurrent falls, particularly among the elderly, should be assessed for underlying BPPV as one of the potential fall-precipitating diagnoses (Jonsson, et al., 2004).

BPPV may occur in the setting of other CNS disorders. Patients should be questioned as to the presence of preexisting CNS disorders that may modify the management of BPPV. BPPV may occur relatively commonly after trauma or traumatic brain injury (Katsarkas, et al., 1999) (Motin, et al., 2005). Posttraumatic BPPV is most likely to involve the posterior semicircular canal, and studies indicate that posttraumatic BPPV is significantly more likely to require repeated physical treatments (up to 67% of cases) for resolution compared with nontraumatic forms (14% of cases) (Gordon, et al., 2004). In rare instances, posttraumatic BPPV may be bilateral (Katsarkas, et al., 1999). Because posttraumatic BPPV may be more refractory and/or bilateral, thus requiring specialized treatment, a history of head trauma preceding a clinical diagnosis of BPPV should be elicited (Motin, et al., 2005). Although dizziness in the setting of multiple sclerosis may have a wide variety of etiologies, studies of acute vertigo occurring in multiple sclerosis report that a substantial number of patients may have BPPV with a positive Dix-Hallpike maneuver and successful response to a PRM (Frohman, et al., 2003) (Frohman, et al., 2000). This study suggests that patients with BPPV and an underlying CNS disorder may be successfully diagnosed and treated with conventional methods for BPPV.

Finally, in a small percentage of cases, refractory or persistent BPPV may create difficulties from a psychological and/or social-functional perspective for affected individuals (Gamiz, et al., 2004) (Lopez-Escamez, et al., 2005). Outcomes studies have shown that patients with BPPV exhibit a significant negative quality-of-life impact from the diagnosis compared with the normative population in multiple subscales of the Short Form-36 (Lopez-Escamez, et al., 2005) (Lopez-Escamez, et al., 2003). Patients who have preexisting comorbid conditions may require additional home supervision in the setting of BPPV (Whitney, et al., 2005). This supervision may include counseling about the risk of falling at home or a home safety assessment. In rare cases, patients disabled by BPPV-related vertigo, especially if chronic or refractory, may need home assistance or temporary nursing home placement for their safety.

Verantwoording

Laatst beoordeeld : 01-02-2020

Laatst geautoriseerd : 01-02-2020

Voor de volledige verantwoording, evidence tabellen en eventuele aanverwante producten raadpleegt u de Richtlijndatabase.

Referenties

- Jönsson, R., Sixt, E., Landahl, S., et al. (2004). Prevalence of dizziness and vertigo in an urban elderly population. *J Vestib Res*, 14, 47-52.
- von Brevem, M., Radtke, A., Lezius, F., et al. (2007). Epidemiology of benign paroxysmal positional vertigo: a population based study. *J Neurol Neurosurg Psychiatry*, 78, 710-5.
- Casellini, C.M., Vinik, A.I. (2007). Clinical manifestations and current treatment options for diabetic neuropathies. *Endocr Pract*, 13, 550-66.
- Cohen, H.S., Kimball, K.T., Stewart, M.G. (2004). Benign paroxysmal positional vertigo and comorbid conditions. *ORL J Otorhinolaryngol Relat Spec*, 66, 11-5.
- Frohman, E.M., Kramer, P.D., Dewey, R.B., et al. (2003). Benign paroxysmal positioning vertigo in multiple sclerosis: diagnosis, pathophysiology and therapeutic techniques. *Mult Scler*, 9, 250-5.
- Frohman, E.M., Zhang, H., Dewey, R.B., et al. (2000). Vertigo in MS: utility of positional and particle repositioning maneuvers. *Neurology*, 55, 1566-9.
- Gamiz, M.J., Lopez-Escamez, J.A. (2004). Health-related quality of life in patients over sixty years old with benign paroxysmal positional vertigo. *Gerontology*, 50, 82-6.
- Gazzola, J.M., Gananca, F.F., Aratani, M.C., et al. (2006). Circumstances and consequences of falls in elderly people with vestibular disorder. *Rev Bras Otorrinolaringol (Engl Ed)*, 72, 388-92.
- Gordon, C.R., Levite, R., Joffe, V., et al. (2004). Is posttraumatic benign paroxysmal positional vertigo different from the idiopathic form?. *Arch Neurol*, 61, 1590-3.
- Katsarkas, A. (1999). Benign paroxysmal positional vertigo (BPPV): idiopathic versus post-traumatic. *Acta Otolaryngol*, 119, 745-9.
- Lawson, J., Johnson, I., Bamiou, D.E., et al. (2005). Benign paroxysmal positional vertigo: clinical characteristics of dizzy patients referred to a Falls and Syncope Unit. *QJM*, 98, 357-64.
- Lopez-Escamez, J.A., Gamiz, M.J., Fernandez-Perez, A., et al. (2005). Long-term outcome and health-related quality of life in benign paroxysmal positional vertigo. *Eur Arch Otorhinolaryngol*, 262, 507-11.
- Motin, M., Keren, O., Groswasser, Z., et al. (2005). Benign paroxysmal positional vertigo as the cause of dizziness in patients after severe traumatic brain injury: diagnosis and treatment. *Brain Inj*, 19, 693-7.
- Richardson, J.K. (2002). Factors associated with falls in older patients with diffuse polyneuropathy. *J Am Geriatr Soc*, 50, 1767-73.
- Rubenstein, L.Z. (2006). Falls in older people: epidemiology, risk factors and strategies for prevention. *Age Ageing*, 35, 37-41.
- Rubenstein, L.Z., Powers, C.M., MacLean, C.H. (2001). Quality indicators for the management and prevention of falls and mobility problems in vulnerable elders. *Ann Intern Med*, 135, 686-93.
- Tilling, L.M., Darawil, K., Britton, M. (2006). Falls as a complication of diabetes mellitus in older people. *J Diabetes Complications*, 20, 158-62.
- Vibert, D., Kompis, M., Hausler, R. (2003). Benign paroxysmal positional vertigo in older women may be related to osteoporosis and osteopenia. *Ann Otol Rhinol Laryngol*, 112, 885-9.

De herevaluatie van de behandeling van BPPD

Het onderwerp 'herevaluatie van behandeling van BPPD' wordt uitgewerkt in verschillende modules.

Verantwoording

Laatst beoordeeld : 01-02-2020

Laatst geautoriseerd : 01-02-2020

Voor de volledige verantwoording, evidence tabellen en eventuele aanverwante producten raadpleegt u de Richtlijndatabase.

De noodzaak van herevaluatie van de behandeling van BPPD

Uitgangsvraag

Is het noodzakelijk om de respons op BPPD behandeling te evalueren?

Aanbeveling

Binnen een maand na de behandeling dient het effect van de behandeling geëvalueerd te worden.

Overwegingen

- Voordeel: identificatie van patiënten met aanhoudende klachten die in eerste instantie met observatie behandeld werden en die zouden kunnen profiteren van de repositiemanoeuvre of een hernieuwde manoeuvre moeten ondergaan. Identificatie van patiënten waarbij de diagnose herzien moet worden.
- Nadeel: geen
- Kosten: kosten van herbeoordeling
- Afweging: het voordeel weegt op tegen het nadeel.
- Waarde oordeel: bevestiging van de diagnose en het ondervangen van patiënten die zouden kunnen profiteren van een andere behandeling.
- Rol van de voorkeur van de patiënt: minimaal.

Onderbouwing

Conclusies

Niveau 3	Het niet reageren op behandeling kan betekenen dat de oorspronkelijke diagnose niet goed was en dat er sprake is van centrale pathologie. Blijvende klachten betekenen een blijvend risico op vallen en verzuim.
-----------------	--

Samenvatting literatuur

Patients with BPPV, regardless of initial treatment option rendered, will have variable responses to therapy (Cohen, et al., 2005). The response to therapy may depend on several factors including the accuracy of the diagnosis of BPPV, the duration of symptoms prior to the diagnosis of BPPV, compliance with prescribed therapy, and other factors (Hilton, et al., 2004) (Rupa, et al., 2004). Patients with BPPV should be reassessed within a set time interval after the diagnosis of BPPV for several reasons.

Failure to respond to initial therapy may indicate an initially erroneous diagnosis of BPPV, and one of the major goals of reassessment is to ensure the accuracy of diagnosis of BPPV. As noted, other more serious CNS disorders may mimic BPPV, and these conditions would not be expected to respond to traditional therapies prescribed for BPPV. In cohort studies, the rate of false-positive diagnosis for BPPV subsequently found to be CNS lesions *after failed treatment* (therefore, a highly selected population) with PRM ranges from 1.1 to 3 percent (Rupa, et al., 2004) (Dal, et al., 2000). Thus, persistence of symptoms after initial management requires clinicians to reassess and reevaluate patients for other etiologies of vertigo. Conversely, resolution of BPPV symptoms after initial therapy such as a PRM would corroborate an accurate diagnosis of BPPV.

Patients who are initially treated with vestibular rehabilitation may fail to resolve symptoms owing to multiple factors including poor compliance. In addition, patients who do not respond to initial therapy are likely to remain at risk for falls, decreased quality of life, and other consequences of unresolved BPPV. For these reasons, patients whose symptoms of BPPV fail to resolve should also be identified and classified as initial treatment failures. To define a treatment failure in BPPV, the clinician needs to determine both a failed outcome criterion and an appropriate time interval for assessment of treatment failure. Successful treatment outcomes for interventions for BPPV are traditionally measured in clinical trials by subjective symptom resolution and/or by conversion to a negative Dix-Hallpike test. Almost all treatment trials for BPPV report an outcome measure in the form of the patient's reported symptoms, typically reported among three categorical outcomes: complete resolution of symptoms, improvement, or no improvement/worsening (Hilton, 2004). When included in meta-analyses, treatment responses are typically incorporated as "all or none" for the complete resolution of vertigo (Hilton, et al., 2004) (Woodworth, et al., 2004) (Teixeira, et al., 2006).

Because effective treatment options are available for BPPV that typically render patients symptom free (if treatment is successful), it is logical to use complete symptom resolution as the outcome of choice at the time of reassessment by the clinician. A symptom-based reassessment also allows clinicians to use clinical judgment as to the most appropriate modality for follow-up for individual patients, including telephone communication, electronic communication, or office based reexamination. This symptom-based assessment of treatment resolution should be detailed enough to distinguish patients with truly decreased symptoms related to treatment or patients with minimized symptoms attributable to positional avoidance (who, in fact, may not be treatment successes) from those with true symptom resolution (Woodworth, et al., 2004).

Although conversion to a negative Dix-Hallpike test may have the advantage of being a more objective reassessment than patients' reported symptoms, it also carries the disadvantage of requiring a repeat clinical visit on the part of the patient with associated direct and indirect costs. The Dix- Hallpike test status is commonly reported in therapeutic trials of BPPV. Persistent symptoms of BPPV and other underlying conditions, however, have been reported in the face of negative Dix-Hallpike testing after therapy, potentially making this a less sensitive reassessment tool (Lynn, et al., 1995) (Magliulo, et al., 2005).

Conversely, patients may report an absence of symptoms after therapeutic intervention yet still have a positive Dix-Hallpike test (Cohen, et al., 2005) (Froehling, et al., 2000) (Sherman, et al., 2001). "Subclinical BPPV" has been offered as an explanation for this (Cohen, et al., 2005). Because of the potential discordance between negative Dix-Hallpike conversion and patients' reported symptoms after treatment for BPPV, Dix- Hallpike conversion is not recommended as the primary reassessment criterion in routine clinical practice but may still be used as a secondary outcome measure. There is no widely accepted time interval at which to assess patients for treatment failure. Therapeutic trials in BPPV variably report follow-up assessments for treatment outcomes at 40 hours, 2 weeks, 1 month, and up to 6 months, although the most commonly chosen interval for follow-up assessment of treatment response is within or at 1 month (Hilton, et al., 2004) (Woodworth, et al., 2004) (Teixeira, et al., 2006). Because the natural history of BPPV exhibits a relatively consistent spontaneous rate of resolution with observation alone, a longer time interval between diagnosis and reassessment would allow patients with true BPPV to resolve symptoms spontaneously, likely irrespective of treatment (Sekine, et al., 2006).

Conversely, the choice of an excessively long time interval between diagnosis and reassessment would also allow cases of an erroneous BPPV diagnosis to potentially progress, leading to potential patient harm. In addition, because recurrence of BPPV may occur as early as 3 months after initial treatment, further delaying the time interval for reassessment may erroneously incorporate a recurrent BPPV syndrome (ie, the initial BPPV responded to treatment with a suitable symptom-free interval thereafter, followed by recurrent BPPV) rather than a persistent BPPV syndrome (Nunez, et al., 2000) (Helminski, et al., 2005).

Given that commonly reported rates of spontaneous complete symptom resolution at the 1-month interval for BPPV range from 20 to 80 percent at 1 month, reassessment at 1 month will also better allow for patients to be reconsidered for further interventional treatment to treat unresolved BPPV (Froehling, et al., 2000) (Lynn, et al., 1995) (Yimtae, et al., 2003) (Munoz, et al., 2007) (Sekine, et al., 2006) (von Brevern, et al., 2006). Thus, choosing a reassessment time interval of 1 month after diagnosis allows a relative balance between overly early reassessment (which would force the unnecessary reassessment of patients who would likely resolve with additional time) and unduly delayed reassessment (which would potentially allow harm from an unknown missed diagnosis or relegate patients to an excess time interval of symptomatic suffering from BPPV). One potential problem with a strict time interval for reassessment is that patients may not have been exposed to their initial treatment (vestibular rehabilitation or PRM as opposed to observation, which may begin immediately after diagnosis) within 1 month of diagnosis depending on referral patterns, patient preferences, or waiting lists for specialty evaluation and treatment. This situation is especially true when the diagnosing clinician may not be the same as the treating clinician. Even if a delay occurs between BPPV diagnosis and completion of the initial treatment, clinicians should still reassess patients at 1 month but may choose to reassign a second time interval for reassessment after completion of the initial treatment option.

Verantwoording

Laatst beoordeeld : 01-02-2020

Laatst geautoriseerd : 01-02-2020

Voor de volledige verantwoording, evidence tabellen en eventuele aanverwante producten raadpleegt u de Richtlijndatabase.

Referenties

- Angeli, S.I., Hawley, R., Gomez, O. (2003). Systematic approach to benign paroxysmal positional vertigo in the elderly. *Otolaryngol Head Neck Surg*, 128, 719-25.
- Asprella Libonati, G. (2005). Diagnostic and treatment strategy of lateral semicircular canal canalolithiasis. *Acta Otorhinolaryngol Ital*, 25, 277-83.
- Bergenius, J., Perols, O. (1999). Vestibular neuritis: a follow-up study. *Acta Otolaryngol*, 119, 895-9.
- Bertholon, P., Bronstein, A.M., Davies, R.A., et al. (2002). Positional down beating nystagmus in 50 patients: cerebellar disorders and possible anterior semicircular canalolithiasis. *J Neurol Neurosurg Psychiatry*, 72, 366-72.
- Beynon, G.J., Baguley, D.M., da Cruz, M.J. (2000). Recurrence of symptoms following treatment of posterior semicircular canal benign positional paroxysmal vertigo with a particle repositioning manoeuvre. *J Otolaryngol*, 29, 2-6.
- Black, F.O., Nashner, L.M. (1984). Postural disturbance in patients with benign paroxysmal positional nystagmus. *Ann Otol Rhinol Laryngol*, 93, 595-9.
- Blatt, P.J., Georgakakis, G.A., Herdman, S.J., et al. (2000). The effect of the canalith repositioning maneuver on resolving postural instability in patients with benign paroxysmal positional vertigo. *Am J Otol*, 21, 356-63.

- von Brevern, M., Seelig, T., Radtke, A., et al. (2006). Short-term efficacy of Epley's manoeuvre: a double-blind randomised trial. *J Neurol Neurosurg Psychiatry*, 77, 980-2.
- Brocchetti, F., Garaventa, G., Ameli, F., et al. (2003). Effect of repetition of Semont's manoeuvre on benign paroxysmal positional vertigo of posterior semicircular canal. *Acta Otorhinolaryngol Ital*, 23, 428-35.
- Buttner, U., Helmchen, C., Brandt, T. (1999). Diagnostic criteria for central versus peripheral positioning nystagmus and vertigo: a review. *Acta Otolaryngol*, 119, 1-5.
- Casani, A.P., Vannucci, G., Fattori, B., et al. (2002). The treatment of horizontal canal positional vertigo: our experience in 66 cases. *Laryngoscope*, 112, 172-8.
- Chang, W.C., Hsu, L.C., Yang, Y.R., et al. (2006). Balance ability in patients with benign paroxysmal positional vertigo. *Otolaryngol Head Neck Surg*, 135, 534-40.
- Chiou, W.Y., Lee, H.L., Tsai, S.C., et al. (2005). A single therapy for all subtypes of horizontal canal positional vertigo. *Laryngoscope*, 115, 1432-5.
- Cohen, H.S., Kimball, K.T. (2005). Effectiveness of treatments for benign paroxysmal positional vertigo of the posterior canal. *Otol Neurotol*, 26, 1034-40.
- Dal, T., Ozluoglu, L.N., Ergin, N.T. (2000). The canalith repositioning maneuver in patients with benign positional vertigo. *Eur Arch Otorhinolaryngol*, 257, 133-6.
- Dornhoffer, J.L., Colvin, G.B. (2000). Benign paroxysmal positional vertigo and canalith repositioning: clinical correlations. *Am J Otol*, 21, 230-3.
- Dunniway, H.M., Welling, D.B. (1998). Intracranial tumors mimicking benign paroxysmal positional vertigo. *Otolaryngol Head Neck Surg*, 118, 429-36.
- Eife, T.D. (1998). Recognition and management of horizontal canal benign positional vertigo. *Am J Otol*, 19, 345-51.
- Froehling, D.A., Bowen, J.M., Mohr, D.N., et al. (2000). The canalith repositioning procedure for the treatment of benign paroxysmal positional vertigo: a randomized controlled trial. *Mayo Clin Proc*, 75, 695-700.
- Furman, J.M., Cass, S.P. (1999). Benign paroxysmal positional vertigo. *N Engl J Med*, 341, 1590-6.
- Furman, J.M., Cass, S.P. (1995). A practical work-up for vertigo. *Contemp Intern Med*, 7, 24-7.
- Giacomini, P.G., Alessandrini, M., Magrini, A. (2002). Long-term postural abnormalities in benign paroxysmal positional vertigo. *ORL J Otorhinolaryngol Relat Spec*, 64, 237-41.
- Gordon, C.R., Levite, R., Joffe, V., et al. (2004). Is posttraumatic benign paroxysmal positional vertigo different from the idiopathic form?. *Arch Neurol*, 61, 1590-3.
- Helminski, J.O., Janssen, I., Kotaspuikis, D., et al. (2005). Strategies to prevent recurrence of benign paroxysmal positional vertigo. *Arch Otolaryngol Head Neck Surg*, 131, 344-8.
- Herdman, S.J., Tusa, R.J. (1996). Complications of the canalith repositioning procedure. *Arch Otolaryngol Head Neck Surg*, 122, 281-6.
- Hilton, M., Pinder, D. (2004). The Epley (canalith repositioning) manoeuvre for benign paroxysmal positional vertigo. *Cochrane Database Syst Rev*, 0, CD003162-.
- Hughes, C.A., Proctor, L. (1997). Benign paroxysmal positional vertigo. *Laryngoscope*, 107, 607-13.
- Jackson, L.E., Morgan, B., Fletcher, J.C., Jr., et al. (2007). Anterior canal benign paroxysmal positional vertigo: an underappreciated entity. *Otol Neurotol*, 28, 218-22.
- Kayan, A., Hood, J.D. (1984). Neuro-otological manifestations of migraine. *Brain*, 107, 1123-42.
- Lynn, S., Lynn, S., Pool, A., Rose, D., et al. (0000). Randomized trial of the canalith repositioning procedure. *Otolaryngol Head Neck Surg*, 113, 712-720.
- Magliulo, G., Bertin, S., Ruggieri, M., et al. (2005). Benign paroxysmal positional vertigo and post-treatment quality of life. *Eur Arch Otorhinolaryngol*, 262, 627-30.
- Monobe, H., Sugawara, K., Murofushi, T. (2001). The outcome of the canalith repositioning procedure for benign paroxysmal positional vertigo: are there any characteristic features of treatment failure cases?. *Acta Otolaryngol Suppl*, 545, 38-40.
- Munoz, J.E., Miklea, J.T., Howard, M., et al. (2007). Canalith repositioning maneuver for benign paroxysmal positional vertigo: randomized controlled trial in family practice. *Can Fam Physician*, 53, 1049-53.
- Nunez, R.A., Cass, S.P., Furman, J.M. (2000). Short- and long-term outcomes of canalith repositioning for benign paroxysmal positional vertigo. *Otolaryngol Head Neck Surg*, 122, 647-52.

- Nuti, D., Agus, G., Barbieri, M.T., et al. (1998). The management of horizontal canal paroxysmal positional vertigo. *Acta Otolaryngol*, 118, 455-60.
- Pollak, L., Davies, R.A., Luxon, L.L. (2002). Effectiveness of the particle repositioning maneuver in benign paroxysmal positional vertigo with and without additional vestibular pathology. *Otol Neurotol*, 23, 79-83.
- Roberts, R.A., Gans, R.E., Kastner, A.H., et al. (2005). Prevalence of vestibulopathy in benign paroxysmal positional vertigo patients with and without prior otologic history. *Int J Audiol*, 44, 191-6.
- Rupa, V. (2004). Persistent vertigo following particle repositioning maneuvers: an analysis of causes. *Arch Otolaryngol Head Neck Surg*, 130, 436-9.
- Sekine, K., Imai, T., Sato, G., et al. (2006). Natural history of benign paroxysmal positional vertigo and efficacy of Epley and Lempert maneuvers. *Otolaryngol Head Neck Surg*, 135, 529-33.
- Sherman, D., Massoud, E.A. (2001). Treatment outcomes of benign paroxysmal positional vertigo. *Journal of Otolaryngology*, 30, 295-9.
- Smouha, E.E., Roussos, C. (1995). Atypical forms of paroxysmal positional nystagmus. *Ear Nose Throat J*, 74, 649-56.
- Teixeira, L.J., Machado, J.N. (2006). Maneuvers for the treatment of benign positional paroxysmal vertigo: a systematic review. *Rev Bras Otorrinolaringol (Engl Ed)*, 72, 130-9.
- Tirelli, G., Russolo, M. (2004). 360-Degree canalith repositioning procedure for the horizontal canal. *Otolaryngol Head Neck Surg*, 131, 740-6.
- Uneri, A. (2004). Migraine and benign paroxysmal positional vertigo: an outcome study of 476 patients. *Ear Nose Throat J*, 83, 814-5.
- White, J.A., Coale, K.D., Catalano, P.J., et al. (2005). Diagnosis and management of lateral semicircular canal benign paroxysmal positional vertigo. *Otolaryngol Head Neck Surg*, 133, 278-84.
- Woodworth, B.A., Gillespie, M.B., Lambert, P.R. (2004). The canalith repositioning procedure for benign positional vertigo: a meta-analysis. *Laryngoscope*, 114, 1143-6.
- Yimtae, K., Srirompotong, S., Sae-Seaw, P. (2003). A randomized trial of the canalith repositioning procedure. *Laryngoscope*, 113, 828-32.

De inhoud van de herevaluatie van de behandeling van BPPD

Uitgangsvraag

Hoe zou de evaluatie van de BPPD behandeling eruit moeten zien?

Aanbeveling

Bij falen van de behandeling moet geëvalueerd worden of de patiënt 1) persistente of andersoortige BPPD heeft 2) co-existente vestibulaire aandoeningen heeft die geïdentificeerd en behandeld moeten worden en 3) een centrale aandoening heeft.

Overwegingen

- Voordeel: het opstellen van een effectief behandelplan voor patiënten met persistente BPPD en comorbiditeiten, afname in de kans op het missen van ernstige aandoeningen die een andere behandeling vereisen.
- Nadeel: geen
- Kosten: kosten van herevaluatie en vervolgtesten
- Afweging: de voordelen wegen op tegen de nadelen.
- Waarde oordeel: behandeling van BPPD en andere vestibulaire aandoeningen die in combinatie kunnen voorkomen; behandeling van persistente BPPD met een repositie manoeuvre na expectatief beleid of vestibulaire revalidatie.
- Rol van de voorkeur van de patiënt: minimaal

Onderbouwing

Conclusies

Niveau 4	Bij falen van de behandeling kan er sprake zijn van 1) persistente of andersoortige BPPD 2) co-existente vestibulaire aandoening 3) centrale aandoening.
-----------------	--

Samenvatting literatuur

Although the repositioning maneuvers have a substantial success rate, they are not expected to solve the at yet unknown origin of clod or debris formation, which may explain the substantial number of re/occurrences of BPPV. Patients with persistent symptoms of vertigo, dizziness, or unsteadiness at the time of reassessment of the initial treatment response are classified as treatment failures. Treatment failures require reevaluation for the following reasons: 1) Persistent BPPV may be present and responsive to additional maneuvers; 2) coexisting vestibular conditions may be present that can be identified and treated; and 3) serious CNS disorders may simulate BPPV and need to be identified (Furman, et al., 1995) (Furman, et al., 1999) (Rupa, et al., 2004).

Persistent BPPV

Patients with BPPV who initially are treated with observation may fail to resolve spontaneously and have persistent BPPV at the time reassessment. Also, on the basis of failure rates of vestibular rehabilitation or a

single-session PRM ranging from 15 to 50 percent, a significant number of patients initially managed with vestibular rehabilitation or PRM will have persistent BPPV at reassessment, which also indicates a treatment failure Furman, et al., 1999) (Hilton, et al., 2004) (Cohen, et al., 2005) (Teixeira, et al., 2006) (von Brevern, et al., 2006). Reevaluation of a treatment failure should include obtaining a history of vertigo, determining if the vertigo is provoked by positional change relative to gravity (ie, lying down in bed, rolling over, bending down, or tilting the head back), which then suggests persistent BPPV. As with the original diagnostic criteria, the Dix-Hallpike test should be repeated to confirm the diagnosis of BPPV. If the Dix-Hallpike maneuver is still positive, repeat PRMs can then be performed as a preferred treatment. The rate of successful treatment of BPPV reaches 90 to 98 percent when additional repositioning maneuvers are subsequently performed (Brochetti, et al., 2003) (Beynon, et al., 2000). Therefore, the PRMs are the treatment of choice for initial BPPV treatment failures deemed to be due to persistent BPPV.

A similar approach may be adopted for the reevaluation of persistent symptoms of vertigo after an initial diagnosis of lateral canal BPPV. The supine roll test should be repeated and, if characteristic nystagmus is elicited, a PRM appropriate for lateral canal BPPV may be repeated as well. There are limited data regarding the management of treatment failures after PRM for lateral canal BPPV, because this condition seems to respond more consistently to PRM and it also has a higher spontaneous resolution rate (Tirelli, et al., 2004) (Sekine, et al., 2006) (Fife, et al., 1998) (Asprella Libonati, et al., 2005). Some studies indicate cure rates of 86 to 100 percent with up to four PRM treatments in lateral canal BPPV (Casani, et al., 2002) (Chiou, et al., 2005). Further subanalysis suggests that the apogeotropic variant of lateral canal BPPV may be more refractory to therapy (White, et al., 2005) (Casani, et al., 2002).

A small percentage of patients initially diagnosed and treated for lateral canal BPPV or horizontal canal BPPV may experience a canal switch. In these cases, initial horizontal canal BPPV may transform into posterior canal BPPV in up to 6 percent of cases (Nutti, et al., 1998) (Tirelli, et al., 2004). Similarly, a small fraction of patients (also approximating 6%) initially presenting with posterior canal BPPV may transition after treatment to lateral canal BPPV (Yimtae, et al., 2003) (Herdman, et al., 1996). A small subset of patients who do not respond to treatment for posterior canal and/or lateral canal BPPV may suffer from anterior canal BPPV, and may need to be evaluated accordingly (Jackson, et al., 2007). Finally, although rare, two semicircular canals may be simultaneously involved. The second canal's involvement may become evident at the time of reassessment if one of the involved canals was appropriately treated (Rupa, et al., 2004). Thus, reassessment of persistent positional vertigo in BPPV should include examination for involvement of other semicircular canals than originally diagnosed. Coexisting Vestibular System Dysfunction A BPPV treatment failure subsequently may be found to be a case manifesting vertiginous symptoms that are provoked by head and body movements in general (ie, not primarily provoked by positional changes relative to gravity); unprovoked (ie, spontaneous) episodes of vertigo occurring while not moving; or in fact, a constant unsteadiness. These specific findings should be identified by clinicians at the time of reevaluation; such findings suggest the presence of vestibular system dysfunction associated with or in addition to the initially treated BPPV. There may be several possible factors at play when vestibular system dysfunction accompanies BPPV.

In a study by Monobe et al, (Monobe, et al., 2001) treatment failure of the PRM was most commonly seen in patients with BPPV secondary to head trauma or vestibular neuritis. Because vestibular neuritis and head trauma are both frequently associated with vestibular dysfunction, the cause of persistent symptoms following

treatment of BPPV is likely related to widespread dysfunction within the vestibular system in this setting (Bergenius, et al., 1999). Because BPPV is more common in patients with Ménière's disease and migraine, vestibular system dysfunction associated with these disorders can lead to prolonged symptoms of BPPV, greater chance for recurrence of BPPV, and increased risk for falls, particularly in older persons (Gordon, et al., 2004) (Roberts, et al., 2005) (Hughes, et al., 1997) (Dornhoffer, et al., 2000) (Uneri, et al., 2004) (Kayan, et al., 1984).

In addition, BPPV not associated with any other ontological or neurological disease can still be associated with an underlying impaired vestibular function, and these individuals are more likely to have incomplete resolution of symptoms even if their Dix-Hallpike testing normalizes with PRM (Pollak, et al., 2002). Finally, transient vestibular dysfunction can also occur following repositioning maneuvers. Evidence suggests that balance function continues to be affected between 1 to 3 months after repositioning maneuvers, and that some of these patients may need additional balance therapy (ie, counseling, vestibular rehabilitation) to prevent falls and decrease their fear of falling after the vertigo from BPPV has resolved (Blatt, et al., 2000) (Chang, et al., 2006) (Giacomini, et al., 2002) (Black, et al., 1984). Thus, reevaluation of BPPV treatment failures should include a search for these associated conditions.

When coexisting vestibular system dysfunction is suspected, additional testing should be considered. This testing may include audiometric testing to screen for Ménière's disease and nerve VIII pathology such as acoustic neuroma, vestibular function testing to detect central and peripheral vestibular dysfunction, and CNS imaging to detect CNS pathology. Such subsequent testing will need to be tailored to the clinical presentation, and clinicians should exercise their clinical judgment. Vestibular rehabilitation has been shown to be an effective treatment for vestibular symptoms due to the potentially persistent vestibular dysfunction associated with BPPV; this treatment may reduce the risk for falls (Angeli, et al., 2003).

CNS Disorders Masquerading as BPPV Although vertigo of central origin is frequently associated with neurological symptoms such as gait, speech, and autonomic dysfunction, it is important to recognize that, rarely, CNS disorders can masquerade as BPPV (Bertholon, et al., 2002). Many of these have been previously discussed in the section on differential diagnosis, but the relative likelihood of their diagnosis increases in the face of initial treatment failure. In one study, a CNS disorder that explained BPPV treatment failure was found in 3 percent of patients (Dal, et al., 2000).

Whenever the signs and symptoms of BPPV are atypical or refractory to treatment, additional history and physical examination should be obtained to address the possibility of undiagnosed CNS disease (Smouha, et al., 1995). Patients with symptoms consistent with those of BPPV who do not show improvement or resolution after undergoing the PRM, especially after two or three attempted maneuvers, or those who describe associated auditory or neurological symptoms should be evaluated with a thorough neurological examination, additional CNS testing, and/or MRI of the brain and posterior fossa to identify possible intracranial pathological conditions (Dunniway, et al., 1998) (Buttner, et al., 1999).

Verantwoording

Laatst beoordeeld : 01-02-2020

Laatst geautoriseerd : 01-02-2020

Voor de volledige verantwoording, evidence tabellen en eventuele aanverwante producten raadpleegt u de Richtlijndatabase.

Referenties

- Angeli, S.I., Hawley, R., Gomez, O. (2003). Systematic approach to benign paroxysmal positional vertigo in the elderly. *Otolaryngol Head Neck Surg*, 128, 719-25.
- Asprella Libonati, G. (2005). Diagnostic and treatment strategy of lateral semicircular canal canalolithiasis. *Acta Otorhinolaryngol Ital*, 25, 277-83.
- Bergenius, J., Perols, O. (1999). Vestibular neuritis: a follow-up study. *Acta Otolaryngol*, 119, 895-9.
- Bertholon, P., Bronstein, A.M., Davies, R.A., et al. (2002). Positional down beating nystagmus in 50 patients: cerebellar disorders and possible anterior semicircular canalolithiasis. *J Neurol Neurosurg Psychiatry*, 72, 366-72.
- Beynon, G.J., Baguley, D.M., da Cruz, M.J. (2000). Recurrence of symptoms following treatment of posterior semicircular canal benign positional paroxysmal vertigo with a particle repositioning manoeuvre. *J Otolaryngol*, 29, 2-6.
- Black, F.O., Nashner, L.M. (1984). Postural disturbance in patients with benign paroxysmal positional nystagmus. *Ann Otol Rhinol Laryngol*, 93, 595-9.
- Blatt, P.J., Georgakakis, G.A., Herdman, S.J., et al. (2000). The effect of the canalith repositioning maneuver on resolving postural instability in patients with benign paroxysmal positional vertigo. *Am J Otol*, 21, 356-63.
- von Brevern, M., Seelig, T., Radtke, A., et al. (2006). Short-term efficacy of Epley's manoeuvre: a double-blind randomised trial. *J Neurol Neurosurg Psychiatry*, 77, 980-2.
- Brocchetti, F., Garaventa, G., Ameli, F., et al. (2003). Effect of repetition of Semont's manoeuvre on benign paroxysmal positional vertigo of posterior semicircular canal. *Acta Otorhinolaryngol Ital*, 23, 428-35.
- Buttner, U., Helmchen, C., Brandt, T. (1999). Diagnostic criteria for central versus peripheral positioning nystagmus and vertigo: a review. *Acta Otolaryngol*, 119, 1-5.
- Casani, A.P., Vannucci, G., Fattori, B., et al. (2002). The treatment of horizontal canal positional vertigo: our experience in 66 cases. *Laryngoscope*, 112, 172-8.
- Chang, W.C., Hsu, L.C., Yang, Y.R., et al. (2006). Balance ability in patients with benign paroxysmal positional vertigo. *Otolaryngol Head Neck Surg*, 135, 534-40.
- Chiou, W.Y., Lee, H.L., Tsai, S.C., et al. (2005). A single therapy for all subtypes of horizontal canal positional vertigo. *Laryngoscope*, 115, 1432-5.
- Cohen, H.S., Kimball, K.T. (2005). Effectiveness of treatments for benign paroxysmal positional vertigo of the posterior canal. *Otol Neurotol*, 26, 1034-40.
- Dal, T., Ozluoglu, L.N., Ergin, N.T. (2000). The canalith repositioning maneuver in patients with benign positional vertigo. *Eur Arch Otorhinolaryngol*, 257, 133-6.
- Dornhoffer, J.L., Colvin, G.B. (2000). Benign paroxysmal positional vertigo and canalith repositioning: clinical correlations. *Am J Otol*, 21, 230-3.
- Dunniway, H.M., Welling, D.B. (1998). Intracranial tumors mimicking benign paroxysmal positional vertigo. *Otolaryngol Head Neck Surg*, 118, 429-36.
- Fife, T.D. (1998). Recognition and management of horizontal canal benign positional vertigo. *Am J Otol*, 19, 345-51.
- Froehling, D.A., Bowen, J.M., Mohr, D.N., et al. (2000). The canalith repositioning procedure for the treatment of benign paroxysmal positional vertigo: a randomized controlled trial. *Mayo Clin Proc*, 75, 695-700.
- Furman, J.M., Cass, S.P. (1999). Benign paroxysmal positional vertigo. *N Engl J Med*, 341, 1590-6.
- Furman, J.M., Cass, S.P. (1995). A practical work-up for vertigo. *Contemp Intern Med*, 7, 24-7.
- Giacomini, P.G., Alessandrini, M., Magrini, A. (2002). Long-term postural abnormalities in benign paroxysmal positional vertigo. *ORL J Otorhinolaryngol Relat Spec*, 64, 237-41.
- Gordon, C.R., Levite, R., Joffe, V., et al. (2004). Is posttraumatic benign paroxysmal positional vertigo different from the idiopathic form?. *Arch Neurol*, 61, 1590-3.

- Helminski, J.O., Janssen, I., Kotaspuikis, D., et al. (2005). Strategies to prevent recurrence of benign paroxysmal positional vertigo. *Arch Otolaryngol Head Neck Surg*, 131, 344-8.
- Herdman, S.J., Tusa, R.J. (1996). Complications of the canalith repositioning procedure. *Arch Otolaryngol Head Neck Surg*, 122, 281-6.
- Hilton, M., Pinder, D. (2004). The Epley (canalith repositioning) manoeuvre for benign paroxysmal positional vertigo. *Cochrane Database Syst Rev*, 0, CD003162-.
- Hughes, C.A., Proctor, L. (1997). Benign paroxysmal positional vertigo. *Laryngoscope*, 107, 607-13.
- Jackson, L.E., Morgan, B., Fletcher, J.C., Jr., et al. (2007). Anterior canal benign paroxysmal positional vertigo: an underappreciated entity. *Otol Neurotol*, 28, 218-22.
- Kayan, A., Hood, J.D. (1984). Neuro-otological manifestations of migraine. *Brain*, 107, 1123-42.
- Lynn, S., Lynn, S., Pool, A., Rose, D., et al. (0000). Randomized trial of the canalith repositioning procedure. *Otolaryngol Head Neck Surg*, 113, 712-720.
- Magliulo, G., Bertin, S., Ruggieri, M., et al. (2005). Benign paroxysmal positional vertigo and post-treatment quality of life. *Eur Arch Otorhinolaryngol*, 262, 627-30.
- Monobe, H., Sugawara, K., Murofushi, T. (2001). The outcome of the canalith repositioning procedure for benign paroxysmal positional vertigo: are there any characteristic features of treatment failure cases?. *Acta Otolaryngol Suppl*, 545, 38-40.
- Munoz, J.E., Miklea, J.T., Howard, M., et al. (2007). Canalith repositioning maneuver for benign paroxysmal positional vertigo: randomized controlled trial in family practice. *Can Fam Physician*, 53, 1049-53.
- Nunez, R.A., Cass, S.P., Furman, J.M. (2000). Short- and long-term outcomes of canalith repositioning for benign paroxysmal positional vertigo. *Otolaryngol Head Neck Surg*, 122, 647-52.
- Nuti, D., Agus, G., Barbieri, M.T., et al. (1998). The management of horizontal canal paroxysmal positional vertigo. *Acta Otolaryngol*, 118, 455-60.
- Pollak, L., Davies, R.A., Luxon, L.L. (2002). Effectiveness of the particle repositioning maneuver in benign paroxysmal positional vertigo with and without additional vestibular pathology. *Otol Neurotol*, 23, 79-83.
- Roberts, R.A., Gans, R.E., Kastner, A.H., et al. (2005). Prevalence of vestibulopathy in benign paroxysmal positional vertigo patients with and without prior otologic history. *Int J Audiol*, 44, 191-6.
- Rupa, V. (2004). Persistent vertigo following particle repositioning maneuvers: an analysis of causes. *Arch Otolaryngol Head Neck Surg*, 130, 436-9.
- Sekine, K., Imai, T., Sato, G., et al. (2006). Natural history of benign paroxysmal positional vertigo and efficacy of Epley and Lempert maneuvers. *Otolaryngol Head Neck Surg*, 135, 529-33.
- Sherman, D., Massoud, E.A. (2001). Treatment outcomes of benign paroxysmal positional vertigo. *Journal of Otolaryngology*, 30, 295-9.
- Smouha, E.E., Roussos, C. (1995). Atypical forms of paroxysmal positional nystagmus. *Ear Nose Throat J*, 74, 649-56.
- Teixeira, L.J., Machado, J.N. (2006). Maneuvers for the treatment of benign positional paroxysmal vertigo: a systematic review. *Rev Bras Otorrinolaringol (Engl Ed)*, 72, 130-9.
- Tirelli, G., Russolo, M. (2004). 360-Degree canalith repositioning procedure for the horizontal canal. *Otolaryngol Head Neck Surg*, 131, 740-6.
- Uneri, A. (2004). Migraine and benign paroxysmal positional vertigo: an outcome study of 476 patients. *Ear Nose Throat J*, 83, 814-5.
- White, J.A., Coale, K.D., Catalano, P.J., et al. (2005). Diagnosis and management of lateral semicircular canal benign paroxysmal positional vertigo. *Otolaryngol Head Neck Surg*, 133, 278-84.
- Woodworth, B.A., Gillespie, M.B., Lambert, P.R. (2004). The canalith repositioning procedure for benign positional vertigo: a meta-analysis. *Laryngoscope*, 114, 1143-6.
- Yimtae, K., Srirompotong, S., Sae-Seaw, P. (2003). A randomized trial of the canalith repositioning procedure. *Laryngoscope*, 113, 828-32.

De voorlichting van patiënten over BPPD

Uitgangsvraag

Ten aanzien van welke aspecten rond BPPD zouden patiënten voorgelicht moeten worden?

Aanbeveling

De patiënt dient verteld te worden dat BPPD een goedaardige, meestal goed behandelbare aandoening is. Het is belangrijk dat de patiënt (eventueel ook de familie) goed voorgelicht wordt over de risico's op vallen bij BPPD, de aanzienlijke kans op een recidief BPPD en het belang van controle door de behandelaar.

Overwegingen

- Voordelen: toename van bewustwording van valrisico, afname van verwondingen door vallen, toename van bewustwording bij patiënten van het risico op terugkeer van BPPD, waardoor ze snel opnieuw voor behandeling komen.
- Nadelen: geen
- Kosten: geen
- Afweging: De voordelen wegen op tegen de nadelen.
- Rol van de voorkeur van de patiënt: geen

Onderbouwing

Conclusies

Niveau 2/3?	Er is bij BPPD sprake van een verhoogde valkans alsook angst om te vallen. Er is tevens een aanzienlijke kans op recidief BPPD na behandeling.
------------------------	---

Samenvatting literatuur

Although BPPV generally responds well to treatment, there is a significant rate of BPPV recurrence after initial resolution or clinical cure. Most trials of BPPV maintain limited follow-up, rarely beyond 3 months. In the few trials of BPPV with longer-term follow-up, the rate of recurrent BPPV (ie, BPPV symptoms manifesting again after a symptom-free period) is reported to be 5 to 13.5 percent at 6-month follow-up (Macias, et al., 2000) (Sridhar, et al., 2005). At 1 year after treatment, the rate of recurrence has been reported at a slightly higher rate of 10 to 18 percent (Prokopakis, et al., 2005) (Sakaida, et al., 2003). The recurrence rate continues to increase and may be as high as 37 to 50 percent at 5 years by Kaplan-Meier estimation (Nunez, et al., 2000) (Sakaida, et al., 2003). Overall the recurrence rate of BPPV may be estimated at 15 percent per year (Nunez, et al., 2000). Patients with BPPV after trauma are likely to demonstrate an even higher recurrence rate of their BPPV (Gordon, et al., 2004). Thus, clinicians should be aware of the recurrence risk of BPPV and should counsel patients accordingly. Counseling will likely have several benefits, which include earlier recognition by patients of recurrent BPPV, allowing earlier return for PRM or vestibular rehabilitation. Also, counseling regarding recurrence will offset the potential anxiety patients may feel when BPPV recurs and allow them to make corresponding adjustments in their daily routine to minimize the impact of BPPV symptomatology. As with any balance or vestibular disorder, patients with BPPV should be counseled regarding the potential that BPPV may place them

at greater risk for falls (Brandt, et al., 1993). This risk may apply particularly to patients with preexisting balance disorders or vestibular deficits and a separate onset of BPPV. The propensity for falling may actually be a significant motivating factor for patients to be referred for evaluation of underlying BPPV (Lawson, et al., 2005). The risk of falls and fear of falls are significant considerations in the management of the elderly who suffer from chronic dizziness (Gazzola, et al., 2006). In a study of 120 elderly patients with chronic vestibular disorders, 36.7 percent carried the diagnosis of BPPV. Fifty-three percent of subjects had fallen at least once in the past year, and 29.2 percent had recurrent falls (Gazzola, et al., 2006). Other authors have confirmed a relatively high rate of BPPV and associated falling tendencies in the elderly (Oghalai, et al., 2000) (Imbaud Genieys, et al., 2007).

Practically speaking, clinicians should counsel patients and their families regarding the risk of falls associated with BPPV. This information is particularly important for the elderly and frail who may be more susceptible to serious injury as a result of falling. Such counseling could include assessment of home safety, activity restrictions, and the need for home supervision until BPPV is resolved (Rubenstein, et al., 2006). Patients may be particularly vulnerable in the time interval between initial diagnosis of BPPV and definitive treatment when they are referred to another clinician for PRM or vestibular rehabilitation. Counseling should therefore occur at the time of initial diagnosis. The costs of such counseling are anticipated to be minimal and will enhance patient and public safety while avoiding potential posttraumatic sequelae.

Finally, patients should be counseled regarding the importance of follow-up after diagnosis of BPPV. Patients initially treated with observation should be counseled that, if BPPV fails to resolve spontaneously, effective therapies such as the PRM may then be undertaken. Also, patients should be educated about atypical symptoms (subjective hearing loss, tinnitus, aural fullness, gait disturbance, non-positional vertigo, nausea, vomiting, etc.) whose occurrence or persistence after resolution of the primary symptoms of BPPV warrant further clinical evaluation (Rupa, et al., 2004). As noted, such symptoms, particularly when unmasked by the resolution of BPPV may indicate an underlying vestibular or CNS disorder. Clinicians may also educate patients with refractory BPPV or repeatedly recurrent BPPV that in select cases a surgical remedy ("canal plugging procedure" or singular neurectomy) may be considered (Parnes, et al., 2003) (Shaia, et al., 2006).

Verantwoording

Laatst beoordeeld : 01-02-2020

Laatst geautoriseerd : 01-02-2020

Voor de volledige verantwoording, evidence tabellen en eventuele aanverwante producten raadpleegt u de Richtlijndatabase.

Referenties

Brandt, T., Dieterich, M. (1993). Vestibular falls. *J Vestib Res*, 3, 3-14.

Gazzola, J.M., Gananca, F.F., Aratani, M.C., et al. (2006). Clinical evaluation of elderly people with chronic vestibular disorder. *Rev Bras Otorrinolaringol (Engl Ed)*, 72, 515-22.

Gordon, C.R., Levite, R., Joffe, V., et al. (2004). Is posttraumatic benign paroxysmal positional vertigo different from the idiopathic form?. *Arch Neurol*, 61, 1590-3.

Imbaud Genieys, S. (2007). Vertigo, dizziness and falls in the elderly. *Annales d Oto-Laryngologie et de Chirurgie Cervico-Faciale*, 124, 189-96.

Macias, J.D., Lambert, K.M., Massingale, S., et al. (2000). Variables affecting treatment in benign paroxysmal positional

vertigo. *Laryngoscope*, 110, 1921-4.

Sridhar, S., Panda, N. (2005). Particle repositioning manoeuvre in benign paroxysmal positional vertigo: is it really safe?. *J Otolaryngol*, 34, 41-5.

Nunez, R.A., Cass, S.P., Furman, J.M. (2000). Short- and long-term outcomes of canalith repositioning for benign paroxysmal positional vertigo. *Otolaryngol Head Neck Surg*, 122, 647-52.

Oghalai, J.S., Manolidis, S., Barth, J.L., et al. (2000). Unrecognized benign paroxysmal positional vertigo in elderly patients. *Otolaryngol Head Neck Surg*, 122, 630-4.

Parnes, L.S., Agrawal, S.K., Atlas, J. (2003). Diagnosis and management of benign paroxysmal positional vertigo (BPPV). *CMAJ*, 169, 681-93.

Prokopakis, E.P., Chimona, T., Tsagournisakis, M., et al. (2005). Benign paroxysmal positional vertigo: 10-year experience in treating 592 patients with canalith repositioning procedure. *Laryngoscope*, 115, 1667-71.

Rubenstein, L.Z. (2006). Falls in older people: epidemiology, risk factors and strategies for prevention. *Age Ageing*, 35, 37-41.

Rupa, V. (2004). Persistent vertigo following particle repositioning maneuvers: an analysis of causes. *Arch Otolaryngol Head Neck Surg*, 130, 436-9.

Sakaida, M., Takeuchi, K., Ishinaga, H., et al. (2003). Long-term outcome of benign paroxysmal positional vertigo. *Neurology*, 60, 1532-4.

Shaia, W.T., Zappia, J.J., Bojrab, D.I., et al. (2006). Success of posterior semicircular canal occlusion and application of the dizziness handicap inventory. *Otolaryngol Head Neck Surg*, 134, 424-30.